

CHARACTERIZING GSVD BY SINGULAR VALUE EXPANSION OF LINEAR OPERATORS AND ITS COMPUTATION*

HAIBO LI†

Abstract. The generalized singular value decomposition (GSVD) of a matrix pair $\{A, L\}$ with $A \in \mathbb{R}^{m \times n}$ and $L \in \mathbb{R}^{p \times n}$ generalizes the singular value decomposition (SVD) of a single matrix. In this paper, we provide a new understanding of GSVD from the viewpoint of SVD, based on which we propose a new iterative method for computing nontrivial GSVD components of a large-scale matrix pair. By introducing two linear operators \mathcal{A} and \mathcal{L} induced by $\{A, L\}$ between two finite-dimensional Hilbert spaces and applying the theory of singular value expansion (SVE) for linear compact operators, we show that the GSVD of $\{A, L\}$ is nothing but the SVEs of \mathcal{A} and \mathcal{L} . This result characterizes completely the structure of GSVD for any matrix pair with the same number of columns. As a direct application of this result, we generalize the standard Golub–Kahan bidiagonalization (GKB) that is a basic routine for large-scale SVD computation such that the resulting generalized GKB (gGKB) process can be used to approximate nontrivial extreme GSVD components of $\{A, L\}$, which is named the gGKB_GSVD algorithm. We use the GSVD of $\{A, L\}$ to study several basic properties of gGKB and also provide preliminary results about convergence and accuracy of gGKB_GSVD for GSVD computation. Numerical experiments are presented to demonstrate the effectiveness of this method.

Key words. GSVD, linear operator, singular value expansion, generalized Golub–Kahan bidiagonalization, Krylov subspace

MSC codes. 15A22, 47A05, 65F99

DOI. 10.1137/24M1651150

1. Introduction. The generalized singular value decomposition (GSVD) of a matrix pair is an extension of the singular value decomposition (SVD) of a single matrix. First introduced by Van Loan [54] and further developed by many others [43, 51, 55], now the GSVD has become a standard matrix decomposition [8, 20]. The GSVD provides an important mathematical tool for analyzing relationships between two sets of variables or matrices, which is particularly useful in various applications, including signal processing [39, 50], statistics [41, 45], computational biology [1], and many others [7, 18, 24, 29, 31].

Let I_k denote the identity matrix of order k and $\mathbf{0}$ denote the zero matrix or vector with dimensions clarified by the context. For any two matrices with the same number of columns, the general-form GSVD is stated as follows [43].

THEOREM 1.1 (GSVD). *Let $A \in \mathbb{R}^{m \times n}$ and $L \in \mathbb{R}^{p \times n}$ with $\text{rank}((A^\top, L^\top)^\top) = r$. Then the GSVD of $\{A, L\}$ is*

$$(1.1a) \quad A = P_A C_A X^{-1}, \quad L = P_L S_L X^{-1}$$

with

$$(1.1b) \quad C_A = \begin{pmatrix} \Sigma_A & \mathbf{0} \\ r & n-r \end{pmatrix}^m, \quad S_L = \begin{pmatrix} \Sigma_L & \mathbf{0} \\ r & n-r \end{pmatrix}^p$$

*Received by the editors April 1, 2024; accepted for publication (in revised form) by M. W. Berry October 10, 2024; published electronically February 20, 2025.

<https://doi.org/10.1137/24M1651150>

†School of Mathematics and Statistics, The University of Melbourne, Parkville, VIC, 3010 Australia (haibo.li@unimelb.edu.au, <https://macheal.github.io/>).

analyzed and developed in several subsequent works; see, e.g., [25, 26, 47]. As a result of the development of contour integration techniques for eigenvalue problems of finding interior eigenvalues [46, 49], recently a contour integral-based algorithm has been adopted for interior GSVD components computation [38].

Apart from regarding the GSVD as an equivalent CSD or GED, there is very little work on understanding and analyzing GSVD from other perspectives. In [10], the authors studied the GSVD using a variational formulation analogous to that of the SVD, providing a new understanding of the generalized singular vectors. Recently, by treating $(A^\top, L^\top)^\top$ (more precisely, the range space of it) as a point in the real Grassmann manifold $\text{Gr}(m+p, r)$ —the manifold composed of all r -dimensional subspaces of \mathbb{R}^{m+p} —the authors in [15] interpreted a modified form of GSVD as a coordinate representation of $(A^\top, L^\top)^\top$. For developing practical GSVD algorithms, however, these new perspectives on GSVD are not enough. It would be beneficial to understand the GSVD from the viewpoint of SVD so that many existing algorithms for large-scale SVD can be adapted for large-scale GSVD computation. One well-known result is that $\{\gamma_i\}$ are the singular values of AL^\dagger if L has full column rank [56], where “ \dagger ” is the Moore–Penrose pseudoinverse. But when L does not have full column rank, generally the GSVD of $\{A, L\}$ is not related to the SVD of AL^\dagger ; this issue becomes much more complicated for nonregular matrix pairs. In [21], the authors proposed a method to compute the leading GSVD components of $\{A, L\}$ by considering an equivalent SVD problem, but the method requires that L is a banded matrix with full row rank and a weighted pseudoinverse [16] should be computed, which is extremely difficult for general matrix pairs.

In this paper, we provide a new understanding of GSVD from the viewpoint of SVD. This new perspective relies on the theory of singular value expansion (SVE) for linear compact operators [17, section 2.2], which is essentially the SVD if the compact operator is a matrix between two Euclidean spaces. Denote by $\mathcal{R}(\cdot)$ and $\mathcal{N}(\cdot)$ the range space and null space of a matrix, respectively. By defining the positive semidefinite matrix $M = A^\top A + L^\top L$, we first investigate the structure of trivial and nontrivial GSVD components x_i , showing that those trivial $\{x_i\}$ form a basis for $\mathcal{N}(M)$ and any nontrivial x_i belongs to the coset $\bar{x}_i + \mathcal{N}(M)$, where $\bar{x}_i \in \mathcal{R}(M)$ is a nontrivial GSVD component. Then we consider the nontrivial $x_i \in \mathcal{R}(M)$ and other corresponding GSVD components. By introducing a linear operator \mathcal{A} induced by $\{A, L\}$ between two finite-dimensional Hilbert spaces, where a non-Euclidean inner product is used, we show that the SVE of \mathcal{A} is just the nontrivial GSVD components of A , i.e., the first decomposition in (1.1a). Similarly, we introduce a linear operator \mathcal{L} induced by $\{A, L\}$ and show that the SVE of \mathcal{L} is just the nontrivial GSVD components of L . This result reveals that the nontrivial part of the GVSD of $\{A, L\}$ is nothing but the SVEs of the two linear operators induced by $\{A, L\}$. Combined with the trivial GSVD components $\{x_{i_j}\}$, it completely characterizes the structure of GSVD for any matrix pair with the same number of columns.

As a direct application of the above result, we propose a new iterative method for computing several extreme nontrivial GSVD components of $\{A, L\}$. This iterative method is a natural extension of the Golub–Kahan bidiagonalization (GKB) method for large-scale SVD computation [19], which iteratively reduces a matrix to a bidiagonal form by a Lanczos-type iterative process. There are several variants and extensions for the standard GKB of a single matrix, which are proposed to solve large-scale generalized least squares problems [3, 6], saddle point problems [14], or regularization of inverse problems [11, 12, 33, 35, 36]; most of them are named the *generalized Golub–Kahan bidiagonalization* (gGKB) while some have other different

names. As a natural analogy of the standard GKB for SVD computation, we develop an operator-type GKB for linear operators \mathcal{A} and \mathcal{L} to approximate their SVE components, which is also named the gGKB process. We derive matrix-form recursive relations for this operator-type GKB so that it can be used in practical computations. Moreover, this approach offers a unified and general treatment for extending the standard GKB, which can be used to derive nearly all of the aforementioned gGKB processes.

Using the GSVD of $\{A, L\}$, we study several basic properties of the proposed gGKB process. Due to the correspondence of GSVD and SVE, the gGKB method can approximate well the extreme nontrivial GSVD components of $\{A, L\}$, resulting in the gGKB_GSVD algorithm. We derive a relative residual norm and its sharp upper bound for the computed nontrivial GSVD components, which is a good measure of the approximating accuracy and can be used in a stopping criterion for practical computations. Several preliminary results about the convergence and accuracy of gGKB_GSVD in exact arithmetic are provided, showing the effectiveness of this method. An advantage of gGKB_GSVD is that it can efficiently compute the extreme nontrivial GSVD components of a large-scale matrix pair, regardless of whether the pair is regular, whereas most existing methods require the matrix pair to be regular.

The paper is organized as follows. In section 2, we review several basic properties of the GSVD. In section 3, we introduce two linear operators induced by $\{A, L\}$ to characterize the structure of GSVD by their SVEs. In section 4 we propose the new gGKB process and study its basic properties. In section 5, we propose the gGKB_GSVD algorithm for computing nontrivial GSVD components. Numerical experiments are presented in section 6, and concluding remarks follow in section 7.

2. GSVD, SVD, and Golub–Kahan bidiagonalization. We review several basic properties of the GSVD of $\{A, L\}$ presented in Theorem 1.1. The nontrivial generalized singular values of $\{A, L\}$ in descending order are

$$(2.1) \quad \underbrace{\infty, \dots, \infty}_{q_1}, \quad \underbrace{c_{q_1+1}/s_{q_1+1}, \dots, c_{q_1+q_2}/s_{q_1+q_2}}_{q_2}, \quad \underbrace{0, \dots, 0}_{q_3}.$$

We remark that the three numbers q_1, q_2, q_3 are uniquely determined by the properties of $\{A, L\}$, and some of them may be zero in certain instances. The nontrivial GSVD components are linked by the vector-form relations

$$(2.2) \quad \begin{cases} Ax_i = c_i p_{A,i}, \\ Lx_i = s_i p_{L,i}, \\ s_i A^\top p_{A,i} = c_i L^\top p_{L,i} \end{cases}$$

for $1 \leq i \leq r$. For those trivial GSVD components, it holds that

$$(2.3) \quad Ax_i = \mathbf{0}, \quad Lx_i = \mathbf{0}, \quad A^\top p_{A,i} = \mathbf{0}, \quad L^\top p_{L,i} = \mathbf{0}$$

for $r+1 \leq i \leq n$. Let \mathcal{P}_S be the projection operator onto a subspace S . The following result describes the structure of trivial and nontrivial GSVD components $\{x_i\}$.

PROPOSITION 2.1. *Let $M = A^\top A + L^\top L$ and partition X as $\begin{pmatrix} X_1 & X_2 \\ r & n-r \end{pmatrix}$. Then $\mathcal{R}(X_2) = \mathcal{N}(M)$. Moreover, let*

$$(2.4) \quad \bar{X} = (\bar{X}_1 \quad X_2), \quad \bar{X}_1 = (\mathcal{P}_{\mathcal{R}(M)}x_1, \dots, \mathcal{P}_{\mathcal{R}(M)}x_r).$$

Then it holds that

$$(2.5) \quad A = P_A C_A \bar{X}^{-1}, \quad L = P_L S_L \bar{X}^{-1}.$$

Proof. It is clear that $\mathcal{N}(M) = \mathcal{N}(A) \cap \mathcal{N}(L)$, and using the GSVD of $\{A, L\}$, it is easy to obtain that $\mathcal{N}(A) \cap \mathcal{N}(L) = \mathcal{R}(X_2)$. Thus, we have $\mathcal{R}(X_2) = \mathcal{N}(M)$. Using the relation that $\mathcal{P}_{\mathcal{R}(M)}x_i = x_i - \mathcal{P}_{\mathcal{N}(M)}x_i$, for $1 \leq i \leq r$ we have

$$A\mathcal{P}_{\mathcal{R}(M)}x_i = Ax_i - A\mathcal{P}_{\mathcal{N}(M)}x_i = Ax_i, \quad L\mathcal{P}_{\mathcal{R}(M)}x_i = Lx_i - L\mathcal{P}_{\mathcal{N}(M)}x_i = Lx_i.$$

Using the above two relations, it is easy to verify (2.5). \square

Since $\dim(\mathcal{N}(M)) = n - r = \text{rank}(\{x_i\}_{i=r+1}^n)$, it follows that $\{x_i\}_{i=r+1}^n$ forms a basis for $\mathcal{N}(M)$. Proposition 2.1 also indicates that for any x_i with $1 \leq i \leq r$, $\mathcal{P}_{\mathcal{R}(M)}x_i$ is also an i th generalized singular vector of $\{A, L\}$. Therefore, any nontrivial x_i can be decomposed into two components, one being $\mathcal{P}_{\mathcal{R}(M)}x_i \in \mathcal{R}(M)$ and the other being an arbitrary vector in $\mathcal{N}(M)$, which means that any nontrivial x_i belongs to the coset $\bar{x}_i + \mathcal{N}(M)$, where $\bar{x}_i \in \mathcal{R}(M)$ is the i th nontrivial GSVD component. In particular, we can focus on those nontrivial x_i in $\mathcal{R}(M)$, which results in the new form of GSVD (2.5). In the subsequent part, we always consider this form of GSVD by requiring

$$(2.6) \quad x_i \in \mathcal{R}(M) \quad \text{for } 1 \leq i \leq r.$$

There exists a direct relationship between the SVD and GSVD for a matrix pair with a special property. If L has full column rank, it follows from (1.1) that $r = n$ and $q_1 = 0$, leading to

$$(2.7) \quad AL^\dagger = P_A C_A X^{-1} [(P_L S_L) X^{-1}]^\dagger = P_A C_A X^{-1} X (P_L S_L)^\dagger = P_A (C_A S_L^\dagger) P_L^\top,$$

where we have used the property that $(M_1 M_2)^\dagger = M_2^\dagger M_1^\dagger$ if M_1 has full column rank and M_2 has full row rank. Therefore, the SVD of AL^\dagger is $P_A (C_A S_L^\dagger) P_L^\top$ with the singular values being $\{\gamma_i\}_{i=1}^n$.

For the above case, one can compute the SVD of AL^\dagger to get the GSVD components.¹ For a large-scale matrix, the GKB process is a basic routine for computing a few extreme SVD components. At each iteration, it reduces the matrix to a lower-order bidiagonal matrix and generates two Krylov subspaces. The Rayleigh–Ritz procedure is then exploited to approximate extreme SVD components using the bidiagonal matrix and Krylov subspaces [4].

In the following part of the paper, we characterize the GSVD from the viewpoint of SVD for any matrix pair with the same number of columns. Then we generalize the GKB process so that it can be used to compute extreme GSVD components.

3. Characterizing GSVD by singular value expansion of linear operators. We first discuss linear operators between two finite-dimensional Hilbert spaces. Then we study the GSVD of $\{A, L\}$ using the SVE of linear operators. Note that subsection 3.1 is quite general without requiring $M = A^\top A + L^\top L$.

3.1. Linear operator induced by matrices. Suppose $G \in \mathbb{R}^{m \times m}$ is symmetric positive definite. It is obvious that $\langle u, u' \rangle_G := u^\top G u'$ defines an inner product on \mathbb{R}^m such that $(\mathbb{R}^m, \langle \cdot, \cdot \rangle_G)$ is an m -dimensional Hilbert space. On the other hand, for any symmetric positive semidefinite matrix $M \in \mathbb{R}^{n \times n}$ with $\text{rank}(M) = r$, the bilinear form $\langle v, v' \rangle_M := v^\top M v'$ is not a well-defined inner product on \mathbb{R}^n if $r < n$. In this case, we consider the inner product on the subspace $\mathcal{R}(M)$.

¹For numerical computations, it is not recommended to explicitly form AL^\dagger due to its numerical instability, especially when L is close to column rank-deficient.

LEMMA 3.1. *The bilinear form $\langle v, v' \rangle_M := v^\top M v'$ for any $v, v' \in \mathcal{R}(M)$ is an inner product, and $(\mathcal{R}(M), \langle \cdot, \cdot \rangle_M)$ is an r -dimensional Hilbert space.*

Proof. The statement $\dim(\mathcal{R}(M)) = \text{rank}(M)$ is a basic property. We need to show that $\langle \cdot, \cdot \rangle_M$ is indeed an inner product, i.e., it is a symmetric and positive bilinear form on $\mathcal{R}(M) \times \mathcal{R}(M)$. We only need to prove the positiveness. To see it, let $v \in \mathcal{R}(M)$ such that $\langle v, v \rangle_M = v^\top M v = 0$. It follows that $v \in \mathcal{N}(M)$. Note that $\mathcal{R}(M) \cap \mathcal{N}(M) = \{\mathbf{0}\}$ since M is symmetric, which leads to $v = \mathbf{0}$. \square

Given a matrix $A \in \mathbb{R}^{m \times n}$, define the linear map

$$(3.1) \quad \mathcal{A}: (\mathcal{R}(M), \langle \cdot, \cdot \rangle_M) \rightarrow (\mathbb{R}^m, \langle \cdot, \cdot \rangle_G), \quad v \mapsto Av,$$

where v and Av are column vectors under the canonical bases of \mathbb{R}^n and \mathbb{R}^m . Let $W_r \in \mathbb{R}^{n \times r}$ and $Z \in \mathbb{R}^{m \times m}$ be two matrices whose columns constitute orthonormal bases of $(\mathcal{R}(M), \langle \cdot, \cdot \rangle_M)$ and $(\mathbb{R}^m, \langle \cdot, \cdot \rangle_G)$, respectively, i.e., $W_r^\top M W_r = I_r$ and $Z^\top G Z = I_m$. Then we have the commutative diagram:

$$(3.2) \quad \begin{array}{ccc} (\mathcal{R}(M), \langle \cdot, \cdot \rangle_M) & \xrightarrow{\mathcal{A}} & (\mathbb{R}^m, \langle \cdot, \cdot \rangle_G) \\ \pi_1 \uparrow & & \uparrow \pi_2 \\ (\mathbb{R}^r, \langle \cdot, \cdot \rangle_2) & \xrightarrow{[\mathcal{A}]} & (\mathbb{R}^m, \langle \cdot, \cdot \rangle_2), \end{array}$$

where $[\mathcal{A}]$ denotes the matrix representation of \mathcal{A} under bases W_r and Z , and π_1 and π_2 are two linear maps:

$$(3.3) \quad \pi_1: (\mathbb{R}^r, \langle \cdot, \cdot \rangle_2) \rightarrow (\mathcal{R}(M), \langle \cdot, \cdot \rangle_M), \quad x \mapsto W_r x,$$

$$(3.4) \quad \pi_2: (\mathbb{R}^m, \langle \cdot, \cdot \rangle_2) \rightarrow (\mathbb{R}^m, \langle \cdot, \cdot \rangle_G), \quad y \mapsto Z y.$$

Note that π_1 and π_2 are two isomorphism maps such that $(\mathcal{R}(M), \langle \cdot, \cdot \rangle_M) \cong (\mathbb{R}^r, \langle \cdot, \cdot \rangle_2)$ and $(\mathbb{R}^m, \langle \cdot, \cdot \rangle_G) \cong (\mathbb{R}^m, \langle \cdot, \cdot \rangle_2)$. Since \mathcal{A} is a bounded linear operator, we can define its adjoint

$$(3.5) \quad \mathcal{A}^*: (\mathbb{R}^m, \langle \cdot, \cdot \rangle_G) \rightarrow (\mathcal{R}(M), \langle \cdot, \cdot \rangle_M), \quad u \mapsto \mathcal{A}^* u$$

by the relation

$$(3.6) \quad \langle \mathcal{A}v, u \rangle_G = \langle \mathcal{A}^* u, v \rangle_M$$

for any $v \in (\mathcal{R}(M), \langle \cdot, \cdot \rangle_M)$ and $u \in (\mathbb{R}^m, \langle \cdot, \cdot \rangle_G)$. We have the following corresponding commutative diagram:

$$(3.7) \quad \begin{array}{ccc} (\mathcal{R}(M), \langle \cdot, \cdot \rangle_M) & \xleftarrow{\mathcal{A}^*} & (\mathbb{R}^m, \langle \cdot, \cdot \rangle_G) \\ \pi_1 \uparrow & & \uparrow \pi_2 \\ (\mathbb{R}^r, \langle \cdot, \cdot \rangle_2) & \xleftarrow{[\mathcal{A}^*]} & (\mathbb{R}^m, \langle \cdot, \cdot \rangle_2), \end{array}$$

where $[\mathcal{A}^*]$ is the matrix representation of \mathcal{A}^* under bases W_r and Z . The following result gives the matrix-forms of $[\mathcal{A}]$ and $[\mathcal{A}^*]$.

LEMMA 3.2. *The matrix representations of \mathcal{A} and \mathcal{A}^* under bases W_r and Z are*

$$(3.8) \quad [\mathcal{A}] = Z^{-1} A W_r, \quad [\mathcal{A}^*] = W_r^\top A^\top G Z.$$

Proof. By the commutative diagram (3.2), we have $\mathcal{A} \circ \pi_1(x) = \pi_2 \circ [\mathcal{A}](x)$ for any $x \in \mathbb{R}^r$, which is equivalent to $AW_r x = Z[\mathcal{A}]x$. Thus, we have $AW_r = Z[\mathcal{A}]$, leading to $[\mathcal{A}] = Z^{-1}AW_r$. Similarly, by the commutative diagram (3.7), we have

$$\mathcal{A}^* \circ \pi_2(y) = \pi_1 \circ [\mathcal{A}^*](y) \Leftrightarrow \mathcal{A}^* Zy = W_r[\mathcal{A}^*]y$$

for any $y \in \mathbb{R}^m$, which leads to

$$(3.9) \quad \mathcal{A}^* Z = W_r[\mathcal{A}^*].$$

From the definition of \mathcal{A}^* , we have $\langle \mathcal{A} \circ \pi_1(x), \pi_2(y) \rangle_G = \langle \pi_1(x), \mathcal{A}^* \circ \pi_2(y) \rangle_M$ for any $x \in \mathbb{R}^r$ and $y \in \mathbb{R}^m$, which can also be written as

$$(AW_r x)^\top G Zy = (W_r x)^\top M(\mathcal{A}^* Zy) \Leftrightarrow x^\top W_r^\top A^\top G Zy = x^\top W_r^\top M \mathcal{A}^* Zy.$$

Thus, we have

$$(3.10) \quad W_r^\top A^\top G Z = W_r^\top M \mathcal{A}^* Z.$$

Combining (3.9) and (3.10) and using $W_r^\top M W_r = I_r$, we finally obtain

$$[\mathcal{A}^*] = W_r^\top M W_r [\mathcal{A}^*] = W_r^\top M \mathcal{A}^* Z = W_r^\top A^\top G Z.$$

The proof is completed. □

The following result will be used throughout the paper.

LEMMA 3.3. *If $\mathcal{R}(W_r) = \mathcal{R}(M)$ and $W_r^\top M W_r = I_r$, then the Moore–Penrose pseudoinverse of M can be expressed as $M^\dagger = W_r W_r^\top$.*

Proof. Let $\bar{M} = W_r W_r^\top$. We only need to verify the following four identities:

$$\begin{aligned} M \bar{M} M &= M, & (M \bar{M})^\top &= M \bar{M}, \\ \bar{M} M \bar{M} &= \bar{M}, & (\bar{M} M)^\top &= \bar{M} M. \end{aligned}$$

The third identity is the easiest to verify: $\bar{M} M \bar{M} = W_r W_r^\top M W_r W_r^\top = W_r W_r^\top = \bar{M}$. Suppose the compact-form eigenvalue decomposition of M is $M = P_r \Lambda_r P_r^\top$ with $\Lambda_r = \text{diag}(\lambda_1, \dots, \lambda_r)$, where $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $P_r \in \mathbb{R}^{n \times r}$ with 2-orthonormal columns. Since $\mathcal{R}(W_r) = \mathcal{R}(M) = \mathcal{R}(P_r)$, there exists $D \in \mathbb{R}^{r \times r}$ such that $W_r = P_r D$. It follows that $I_r = W_r^\top M W_r = D^\top \Lambda_r D$. Therefore, it follows that $M \bar{M} M = M P_r D D^\top P_r^\top M = P_r \Lambda_r D D^\top \Lambda_r P_r^\top = P_r \Lambda_r P_r^\top = M$, since $\Lambda_r D D^\top = D^\top \Lambda_r D = I_r$. Similarly, we have

$$\begin{aligned} M \bar{M} &= P_r \Lambda_r P_r^\top P_r D D^\top P_r^\top = P_r \Lambda_r D D^\top P_r^\top = P_r P_r^\top = (M \bar{M})^\top, \\ \bar{M} M &= P_r D D^\top P_r^\top M = P_r D D^\top \Lambda_r P_r^\top = P_r P_r^\top = (\bar{M} M)^\top. \end{aligned}$$

Now all the four identities have been verified. □

3.2. Characterizing GSVD by singular value expansion. In this subsection, we consider the simpler case that $G = I_m$, since it has direct connections with the GSVD of a matrix pair. For notational simplicity, let $\mathcal{X} = (\mathcal{R}(M), \langle \cdot, \cdot \rangle_M)$ and $\mathcal{Y} = (\mathbb{R}^m, \langle \cdot, \cdot \rangle_2)$. For the linear compact operator

$$(3.11) \quad \mathcal{A}: \mathcal{X} \rightarrow \mathcal{Y}, \quad v \mapsto Av$$

between the two Hilbert spaces \mathcal{X} and \mathcal{Y} , where v and Av are column vectors under the canonical bases of \mathbb{R}^n and \mathbb{R}^m , it has the SVE with finite terms; see, e.g.,

[30, section 15.4]. Here we use the terminology ‘‘SVE’’ instead of ‘‘SVD’’ to distinguish it from the SVD of a matrix. The theory of SVE for \mathcal{A} states that there exist positive scalars $\sigma_1 \geq \dots \geq \sigma_d > 0$, two orthonormal systems $\{f_i\}_{i=1}^d \subseteq \mathcal{X}$ and $\{h_i\}_{i=1}^d \subseteq \mathcal{Y}$ such that

$$(3.12) \quad \mathcal{A}f_i = \sigma_i h_i, \quad \mathcal{A}^* h_i = \sigma_i f_i,$$

and any $v \in \mathcal{X}$ has the expansion

$$(3.13) \quad v = v_0 + \sum_{i=1}^d \langle v, f_i \rangle_M f_i$$

with some $v_0 \in \ker(\mathcal{A})$, and

$$(3.14) \quad \mathcal{A}v = \sum_{i=1}^d \sigma_i \langle v, f_i \rangle_M h_i,$$

where $d = \dim(\text{im}(\mathcal{A}))$. Here we use $\ker(\cdot)$ and $\text{im}(\cdot)$ to denote the kernel and image of a linear operator, respectively, to distinguish them from the null space $\mathcal{N}(\cdot)$ and range space $\mathcal{R}(\cdot)$ of a matrix.

The following result provides more details about the SVE of \mathcal{A} .

THEOREM 3.4. *For any $A \in \mathbb{R}^{m \times n}$ and symmetric positive semidefinite matrix $M \in \mathbb{R}^{n \times n}$ with rank r , define the linear operator \mathcal{A} as (3.11). Then there exist an M -orthonormal matrix $F \in \mathbb{R}^{n \times r}$, a 2-orthonormal matrix $H \in \mathbb{R}^{m \times m}$, and a diagonal matrix $\Sigma \in \mathbb{R}^{m \times r}$, such that for any $v \in \mathcal{X}$ and $u \in \mathcal{Y}$, it holds that*

$$(3.15) \quad \mathcal{A}v = H\Sigma F^\top Mv, \quad \mathcal{A}^*u = F\Sigma^\top H^\top u$$

under the canonical bases of \mathbb{R}^n and \mathbb{R}^m .

Proof. Let $\mathcal{X}_1 = \text{span}\{f_i\}_{i=1}^d$. We first prove $\mathcal{X} = \ker(\mathcal{A}) \oplus \mathcal{X}_1$. Noticing (3.13), we only need to prove $\ker(\mathcal{A}) \cap \mathcal{X}_1 = \{\mathbf{0}\}$. Let $v = \sum_{j=1}^d \mu_j f_j \in \ker(\mathcal{A}) \cap \mathcal{X}_1$. By (3.14), it follows that $\mathbf{0} = \mathcal{A}v = \sum_{j=1}^d \sigma_j \mu_j h_j$, leading to $\sigma_i \mu_i = 0$ for $1 \leq i \leq d$. Since $\sigma_i > 0$, we have $\mu_i = 0$ for $1 \leq i \leq d$, and thereby $v = \mathbf{0}$. Then we prove $\ker(\mathcal{A}) \perp_M \mathcal{X}_1$, where \perp_M is the orthogonal relation in \mathcal{X} . For any $v \in \ker(\mathcal{A})$ and any f_i , by (3.12) we have $f_i = \sigma_i^{-1} \mathcal{A}^* h_i$, and thereby

$$\langle v, f_i \rangle_M = \langle v, \sigma_i^{-1} \mathcal{A}^* h_i \rangle_M = \langle \mathcal{A}v, \sigma_i^{-1} h_i \rangle_2 = \langle \mathbf{0}, \sigma_i^{-1} h_i \rangle_2 = 0.$$

Note $\dim(\mathcal{X}_1) = d$. Therefore, we can find $r - d$ M -orthonormal vectors in $\ker(\mathcal{A})$ that are M -orthogonal to each f_i . Denote these vectors by $\{f_{d+1}, \dots, f_r\}$. Then $\{f_i\}_{i=1}^r$ constitute a complete orthonormal basis for \mathcal{X} . From (3.14) we have $\text{im}(\mathcal{A}) = \text{span}\{h_i\}_{i=1}^d =: \mathcal{Y}_1$. Using the relation $\ker(\mathcal{A}^*) = \text{im}(\mathcal{A})^\perp = \mathcal{Y}_1^\perp$, where the orthogonality is taken in $(\mathbb{R}^m, \langle \cdot, \cdot \rangle_2)$, there exist $m - d$ 2-orthonormal $\{h_{d+1}, \dots, h_m\} \subseteq \ker(\mathcal{A}^*)$ such that $\{h_i\}_{i=1}^m$ constitute a complete orthonormal basis for \mathcal{Y} .

Therefore, for any $v \in \mathcal{X}$, it can be written as $v = \sum_{i=1}^r \langle v, f_i \rangle_M f_i$, and thereby

$$\mathcal{A}v = \sum_{i=1}^r \langle v, f_i \rangle_M \mathcal{A}f_i = \sum_{i=1}^r \sigma_i \langle v, f_i \rangle_M h_i,$$

where we define $\sigma_{d+1} = \dots = \sigma_r = 0$. Similarly, for any $u \in \mathcal{Y}$ with the expansion $u = \sum_{i=1}^m \langle u, h_i \rangle_2 h_i$, it holds that

$$\mathcal{A}^*u = \sum_{i=1}^m \langle u, h_i \rangle_M \mathcal{A}^*h_i = \sum_{i=1}^r \sigma_i \langle u, h_i \rangle_2 f_i.$$

Let the matrices $F = (f_1, \dots, f_r)$, $H = (h_1, \dots, h_m)$, and $\Sigma = (\Sigma_d \mathbf{0}) \in \mathbb{R}^{m \times r}$ with $\Sigma_d = \text{diag}(\sigma_1, \dots, \sigma_d)$. Then (3.15) is just the matrix-form of the above two identities. \square

One can verify that (3.12)–(3.14) can be derived from (3.15). Therefore, the two relations in (3.15) describe completely the SVE of \mathcal{A} . In the following part, we use the notation

$$(3.16) \quad \mathcal{A} \sim H \Sigma F^\top$$

to denote the SVE of \mathcal{A} . From the proof of Theorem 3.4, we have the following basic relations for the four important subspaces:

$$(3.17a) \quad \ker(\mathcal{A}) = \text{span}\{f_i\}_{i=d+1}^r, \quad \text{im}(\mathcal{A}) = \text{span}\{h_i\}_{i=1}^d,$$

$$(3.17b) \quad \ker(\mathcal{A}^*) = \text{span}\{h_i\}_{i=d+1}^m, \quad \text{im}(\mathcal{A}^*) = \text{span}\{f_i\}_{i=1}^d.$$

From the theory of SVE for linear compact operators, if the multiplicity of σ_i is 1, then the corresponding f_i and h_i are uniquely determined (at most differing by a sign). If the multiplicity of σ_i is $m_i > 1$, then there are m_i corresponding linearly independent $\{f_i\}$ and $\{h_i\}$, respectively, which are M -orthonormal and 2-orthonormal.

Based on Theorem 3.4, now we can use the SVE of \mathcal{A} to characterize the GSVD of $\{A, L\}$. Remember that we consider that $x_i \in \mathcal{R}(M)$ for $1 \leq i \leq r$.

THEOREM 3.5. *Let the GSVD of $\{A, L\}$ be (1.1) and let \mathcal{A} be defined as (3.11) with $M = A^\top A + L^\top L$. Partition P_A and X as*

$$(3.18) \quad P_A = \begin{pmatrix} P_{A1} & P_{A2} & P_{A3} \\ q_1 & q_2 & m - q_1 - q_2 \end{pmatrix} m, \quad X = \begin{pmatrix} X_1 & X_2 & X_3 & X_4 \\ q_1 & q_2 & q_3 & n - r \end{pmatrix} n$$

and let $\tilde{X}_1 = (X_1 \ X_2 \ X_3)$. Then the SVE of \mathcal{A} has the form

$$(3.19) \quad \mathcal{A} \sim P_A \Sigma_A \tilde{X}_1^\top,$$

and it holds that

$$(3.20a) \quad \ker(\mathcal{A}) = \mathcal{R}(X_3), \quad \text{im}(\mathcal{A}) = \mathcal{R}((P_{A1} \ P_{A2})),$$

$$(3.20b) \quad \ker(\mathcal{A}^*) = \mathcal{R}(P_{A3}), \quad \text{im}(\mathcal{A}^*) = \mathcal{R}((X_1 \ X_2)).$$

Proof. Using the GSVD of $\{A, L\}$, we have

$$A^\top A + L^\top L = X^{-\top} \left(\begin{pmatrix} \Sigma_A^\top \Sigma_A & \\ & \mathbf{0} \end{pmatrix} + \begin{pmatrix} \Sigma_L^\top \Sigma_L & \\ & \mathbf{0} \end{pmatrix} \right) X^{-1} = X^{-\top} \begin{pmatrix} I_r & \\ & \mathbf{0} \end{pmatrix} X^{-1},$$

which leads to $\text{rank}(M) = r$ and

$$\begin{pmatrix} I_r & \\ & \mathbf{0} \end{pmatrix} = \begin{pmatrix} \tilde{X}_1^\top \\ X_4^\top \end{pmatrix} M \begin{pmatrix} \tilde{X}_1 & X_4 \end{pmatrix} = \begin{pmatrix} \tilde{X}_1^\top M \tilde{X}_1 & \tilde{X}_1^\top M X_4 \\ X_4^\top M \tilde{X}_1 & X_4^\top M X_4 \end{pmatrix}.$$

Therefore, we have $\tilde{X}_1^\top M \tilde{X}_1 = I_r$. Note that $\mathcal{R}(\tilde{X}_1) \subseteq \mathcal{R}(M)$. It follows that \tilde{X}_1 is an M -orthonormal basis of $(\mathcal{R}(M), \langle \cdot, \cdot \rangle_M)$, and thereby we obtain from Lemma 3.3 that $M^\dagger = \tilde{X}_1 \tilde{X}_1^\top$. Notice from (1.1) that

$$A(\tilde{X}_1 \ X_4) = P_A(\Sigma_A \ \mathbf{0}) \Rightarrow A\tilde{X}_1 = P_A \Sigma_A.$$

Thus, we have $AM^\dagger = A\tilde{X}_1\tilde{X}_1^\top = P_A\Sigma_A\tilde{X}_1^\top$. For any $v \in (\mathcal{R}(M), \langle \cdot, \cdot \rangle_M)$, it holds that

$$\mathcal{A}v = \mathcal{A}\mathcal{P}_{\mathcal{R}(M)}v = AM^\dagger Mv = P_A\Sigma_A\tilde{X}_1^\top Mv.$$

Using the commutative diagram (3.7) and noticing $G = Z = I_m$ for the current case, we have for any $u \in (\mathbb{R}^m, \langle \cdot, \cdot \rangle_2)$ that

$$(3.21) \quad \mathcal{A}^*u = \pi_1 \circ [\mathcal{A}^*](u) = \tilde{X}_1(\tilde{X}_1^\top A^\top)u = M^\dagger A^\top u = (AM^\dagger)^\top u = \tilde{X}_1\Sigma_A^\top P_A^\top u,$$

where we have used $[\mathcal{A}^*] = \tilde{X}_1^\top A^\top$ by Lemma 3.2. This proves that the SVE of \mathcal{A} has the form $P_A\Sigma_A\tilde{X}_1^\top$.

From the SVE of \mathcal{A} we have $\dim(\text{im}(\mathcal{A})) = q_1 + q_2$. Using the relations (3.17), it follows that $\text{im}(\mathcal{A}) = \mathcal{R}((P_{A1} \ P_{A2}))$. Since P_A is a 2-orthogonal matrix, we then have $\ker(\mathcal{A}^*) = \mathcal{R}(P_{A3})$. The other two relations can also be verified easily. \square

Corresponding to Theorem 3.5, we have the following result.

THEOREM 3.6. *Define \mathcal{L} as*

$$(3.22) \quad \mathcal{L} : (\mathcal{R}(M), \langle \cdot, \cdot \rangle_M) \rightarrow (\mathbb{R}^p, \langle \cdot, \cdot \rangle_2), \quad v \mapsto Lv,$$

where v and Lv are column vectors under the canonical bases of \mathbb{R}^n and \mathbb{R}^p . Partition P_L as

$$(3.23) \quad P_L = \begin{pmatrix} P_{L1} & P_{L2} & P_{L3} \\ p - q_2 - q_3 & q_2 & q_3 \end{pmatrix} p.$$

Then the SVE of \mathcal{L} has the form

$$(3.24) \quad \mathcal{L} \sim P_L\Sigma_L\tilde{X}_1^\top,$$

and it holds that

$$(3.25a) \quad \ker(\mathcal{L}) = \mathcal{R}(X_1), \quad \text{im}(\mathcal{L}) = \mathcal{R}((P_{L2} \ P_{L3})),$$

$$(3.25b) \quad \ker(\mathcal{L}^*) = \mathcal{R}(P_{L1}), \quad \text{im}(\mathcal{L}^*) = \mathcal{R}((X_2 \ X_3)).$$

Proposition 2.1 together with Theorems 3.5 and 3.6 characterizes completely the GSVD of $\{A, L\}$ based on the SVEs of linear operators \mathcal{A} and \mathcal{L} . Particularly, they show that the nontrivial part \tilde{X}_1 is the common right SVE components of \mathcal{A} and \mathcal{L} , while P_A and P_L are the left SVE components of \mathcal{A} and \mathcal{L} , respectively. Moreover, the relations (3.20) and (3.25) use the image spaces and kernel spaces of \mathcal{A} , \mathcal{A}^* and \mathcal{L} , \mathcal{L}^* to describe the structure of each GSVD blocks and give a new explanation of the three numbers q_1 , q_2 , and q_3 in (1.1).

Based on the SVE characterization of GSVD, we can expect to modify those algorithms for large-scale SVD computation for large-scale GSVD computation. To compute nontrivial extreme GSVD components, we generalize the standard GKB process from the viewpoint of linear operators.

4. Generalizing the Golub–Kahan bidiagonalization. In this section, the generalization of GKB is quite general without requiring $M = A^\top A + L^\top L$, and we follow the notation and assumptions in subsection 3.1. For the linear operator in (3.1), the iterative process of GKB can be described as follows; see [9] for discussions

about GKB for linear compact operators. Choosing a nonzero vector $b \in (\mathbb{R}^m, \langle \cdot, \cdot \rangle_G)$, the basis recursive relations are

$$(4.1) \quad \begin{cases} \beta_1 u_1 = b, \\ \alpha_i v_i = \mathcal{A}^* u_i - \beta_i v_{i-1}, \\ \beta_{i+1} u_{i+1} = \mathcal{A} v_i - \alpha_i u_i, \end{cases}$$

where $u_i \in (\mathbb{R}^m, \langle \cdot, \cdot \rangle_G)$ and $v_i \in (\mathcal{R}(M), \langle \cdot, \cdot \rangle_M)$, and α_i and β_i are positive scalars such that $\|v_i\|_M = \|u_i\|_G = 1$. Note that $v_0 := \mathbf{0}$ for the initial step.

For the purpose of practical computation, we need to derive a matrix-form expression of the recursive relations. Using the isomorphisms π_1 and π_2 , denote u_i and v_i by $u_i = Z y_i$ and $v_i = W_r x_i$ with $y_i \in \mathbb{R}^m$ and $x_i \in \mathbb{R}^r$ for any $i \geq 1$. Then we have

$$\begin{aligned} \mathcal{A} v_i &= \mathcal{A} \circ \pi_1(x_i) = \pi_2 \circ [\mathcal{A}] x_i = Z Z^{-1} \mathcal{A} W_r x_i = A v_i, \\ \mathcal{A}^* u_i &= \mathcal{A}^* \circ \pi_2(y_i) = \pi_1 \circ [\mathcal{A}^*] y_i = W_r W_r^\top A^\top G Z y_i = M^\dagger A^\top G u_i. \end{aligned}$$

Therefore, the last two recursions in (4.1) can be written in the matrix-vector forms

$$(4.2) \quad \begin{cases} \alpha_i v_i = M^\dagger A^\top G u_i - \beta_i v_{i-1}, \\ \beta_{i+1} u_{i+1} = A v_i - \alpha_i u_i. \end{cases}$$

Using (4.2), the GKB of \mathcal{A} can proceed step by step. We name the above iterative process the **gGKB**. The pseudocode of **gGKB** is shown in Algorithm 4.1.

Remark 4.1. If G is also positive semidefinite, define the linear operator $\mathcal{A} : (\mathcal{R}(M), \langle \cdot, \cdot \rangle_M) \rightarrow (\mathcal{R}(G), \langle \cdot, \cdot \rangle_G)$ to be $v \mapsto Av$. In this case, a similar **gGKB** process can be obtained. A slight difference is that the initial vector should satisfy $b \in \mathcal{R}(G)$.

For large-scale matrices, we cannot directly compute M^\dagger . In this case, using the relation

$$(4.3) \quad M^\dagger \bar{s} = \underset{s \in \mathbb{R}^n}{\operatorname{argmin}} \|M s - \bar{s}\|_2,$$

we can compute $M^\dagger \bar{s}$ by iteratively solving the above least squares problems. If we use **gGKB** to compute the GSVD of $\{A, L\}$ (see section 5), which means $G = I_m$, then $M^\dagger \bar{s} = M^\dagger A^\top u_i$ is the minimum 2-norm solution to the least squares problem

$$(4.4) \quad \min_{s \in \mathbb{R}^n} \left\| \begin{pmatrix} A \\ L \end{pmatrix} s - \begin{pmatrix} u_i \\ \mathbf{0} \end{pmatrix} \right\|_2.$$

Algorithm 4.1. Generalized Golub–Kahan bidiagonalization.

Input: $A \in \mathbb{R}^{m \times n}$, $M \in \mathbb{R}^{n \times n}$, $G \in \mathbb{R}^{m \times m}$, $b \in \mathbb{R}^m$

- 1: Initialize: let $\beta_1 = \|b\|_G$, $u_1 = b/\beta_1$
- 2: Compute $\bar{s} = A^\top G u_1$, $s = M^\dagger \bar{s}$
- 3: $\alpha_1 = \|s\|_M$, $v_1 = s/\alpha_1$
- 4: **for** $i = 1, 2, \dots, k$, **do**
- 5: $q = A v_i - \alpha_i u_i$
- 6: $\beta_{i+1} = \|q\|_G$, $u_{i+1} = q/\beta_{i+1}$
- 7: $\bar{s} = A^\top G u_{i+1}$, $s = M^\dagger \bar{s} - \beta_{i+1} v_i$
- 8: $\alpha_{i+1} = \|s\|_M$, $v_{i+1} = s/\alpha_{i+1}$
- 9: **end for**

Output: $\{\alpha_i, \beta_i\}_{i=1}^{k+1}$, $\{u_i, v_i\}_{i=1}^{k+1}$

Both (4.3) and (4.4) can be iteratively solved by using the LSQR algorithm [44]. If $\begin{pmatrix} A \\ L \end{pmatrix}$ is sparse, then it is more efficient to compute $M^\dagger \bar{s}$ by solving (4.4). For both cases, gGKB has a nested inner-outer iteration structure.

If $G = I_m$ and $M = I_n$, then gGKB becomes the standard GKB. If $G = I_m$ and M is invertible, the gGKB is equivalent to the generalizations of GKB proposed in [3, 11, 12, 33, 35] with different forms. The following result describes the basic property of gGKB, very similar to that of the standard GKB.

PROPOSITION 4.1. *For each v_i generated by gGKB, it holds that $v_i \in \mathcal{R}(M)$. The group of vectors $\{v_i\}_{i=1}^k$ is an M -orthonormal basis of the Krylov subspace*

$$(4.5) \quad \mathcal{K}_k(M^\dagger A^\top G A, M^\dagger A^\top G b) = \text{span}\{(M^\dagger A^\top G A)^i M^\dagger A^\top G b\}_{i=0}^{k-1},$$

and $\{u_i\}_{i=1}^k$ is a G -orthonormal basis of the Krylov subspace

$$(4.6) \quad \mathcal{K}_k(A M^\dagger A^\top G, b) = \text{span}\{(A M^\dagger A^\top G)^i b\}_{i=0}^{k-1}.$$

Proof. To get a better understanding of gGKB, we give two proofs.

The first proof. We prove $v_i \in \mathcal{R}(M)$ by mathematical induction. First note $\mathcal{R}(M^\dagger) = \mathcal{R}(M)$ for a symmetric M . For $i = 1$, we have $\alpha_1 = M^\dagger A^\top G u_1 \in \mathcal{R}(M)$. Suppose $v_i \in \mathcal{R}(M)$ for $i \geq 1$. From the recursion (4.2) we obtain

$$\alpha_{i+1} v_{i+1} = M^\dagger A^\top G u_i - \beta_{i+1} v_i \in \mathcal{R}(M),$$

leading to $v_{i+1} \in \mathcal{R}(M)$. To prove the second property, we exploit the theory about the GKB of $\mathcal{A} : (\mathcal{R}(M), \langle \cdot, \cdot \rangle_M) \rightarrow (\mathbb{R}^m, \langle \cdot, \cdot \rangle_G)$ with starting vector b , which states that $\{v_i\}_{i=1}^k$ and $\{u_i\}_{i=1}^k$ are the M -orthonormal basis and G -orthonormal basis of the two Krylov subspaces:

$$\begin{aligned} \mathcal{K}_k(\mathcal{A}^* \mathcal{A}, \mathcal{A}^* b) &:= \text{span}\{(\mathcal{A}^* \mathcal{A})^i \mathcal{A}^* b\}_{i=0}^{k-1}, \\ \mathcal{K}_k(\mathcal{A} \mathcal{A}^*, b) &:= \text{span}\{(\mathcal{A} \mathcal{A}^*)^i b\}_{i=0}^{k-1}, \end{aligned}$$

respectively. For any $u \in \mathbb{R}^m = \pi_2(y) = Zy$, from the commutative diagram (3.7) and using Lemmas 3.2 and 3.3, we obtain

$$\mathcal{A}^* u = \mathcal{A}^* \circ \pi_2(y) = \pi_1 \circ [\mathcal{A}^*](y) = W_r (W_r^\top A^\top G Z) y = M^\dagger A^\top G u.$$

Therefore, we have

$$(\mathcal{A}^* \mathcal{A})^i \mathcal{A}^* b = (M^\dagger A^\top G A)^i M^\dagger A^\top G b, \quad (\mathcal{A} \mathcal{A}^*)^i b = (A M^\dagger A^\top G)^i b.$$

The second proof. Using the coordinates of u_i and v_i under bases W_r and Z , we can write the last two relations in (4.1) as

$$\alpha_i W_r x_i = \mathcal{A}^* Z y_i - \beta_i W_r x_{i-1}, \quad \beta_{i+1} Z y_{i+1} = \mathcal{A} W_r x_i - \alpha_i Z y_i,$$

where $v_i = W_r x_i$ and $u_i = Z y_i$. Note that $Z^\top G Z = I_m$ implies $Z^{-1} = Z^\top G$. Letting $\bar{b} = Z^{-1} b$, multiplying from left the first and second relations by $W_r^\top M$ and Z^{-1} , and using (3.10), we obtain

$$(4.7) \quad \begin{cases} \beta_1 y_1 = \bar{b}_1, \\ \alpha_i x_i = W_r^\top A^\top G Z y_i - \beta_i x_{i-1}, \\ \beta_{i+1} y_{i+1} = Z^\top G A W_r x_i - \alpha_i y_i. \end{cases}$$

Since $G^\top = G$, it follows that (4.7) is the standard GKB of matrix $Z^\top GAW_r$ with starting vector \bar{b} . Therefore, $\{x_i\}_{i=1}^k$ and $\{y_i\}_{i=1}^k$ are 2-orthonormal bases of the two Krylov subspaces

$$\begin{aligned} & \text{span}\{((Z^\top GAW_r)^\top Z^\top GAW_r)^i (Z^\top GAW_r)^\top \bar{b}\}_{i=0}^{k-1}, \\ & \text{span}\{(Z^\top GAW_r (Z^\top GAW_r)^\top)^i \bar{b}\}_{i=0}^{k-1}, \end{aligned}$$

respectively. Note that

$$\begin{aligned} & W_r((Z^\top GAW_r)^\top Z^\top GAW_r)^i (Z^\top GAW_r)^\top \bar{b} \\ &= W_r(W_r^\top A^\top G Z Z^\top GAW_r)^i W_r^\top A^\top G Z \bar{b} = W_r(W_r^\top A^\top GAW_r)^i W_r^\top A^\top Gb \\ &= (W_r W_r^\top A^\top GA)^i W_r W_r^\top A^\top Gb = (M^\dagger A^\top GA)^i M^\dagger A^\top Gb. \end{aligned}$$

We have $v_i = W_r x_i \in \mathcal{R}(M)$ and obtain (4.5). Similarly, we can obtain (4.6). \square

It is easy to verify that (4.7) is equivalent to (4.1). Note from Lemma 3.2 that $[\mathcal{A}^*] = [A]^\top$ since $Z^{-1} = Z^\top G$. Therefore, the matrix representations of \mathcal{A} and \mathcal{A}^* are $Z^\top GAW_r \in \mathbb{R}^{m \times r}$ and $(Z^\top GAW_r)^\top$, respectively, which maps a coordinate vector from \mathbb{R}^r to \mathbb{R}^m . In this sense, we can say that the recursive relation (4.7) is the coordinate representation for the gGKB of \mathcal{A} under bases W_r and Z .

From the first proof of Proposition 4.1, we have $s \in \mathcal{R}(M)$. Therefore, for each $i \geq 1$, if $s = M^\dagger A^\top Gu_i - \beta_i v_{i-1} \neq \mathbf{0}$, then $\alpha_i = \|s\|_M \neq 0$. This indicates that even if M is not positive definite, the gGKB does not terminate as long as s or $q = Av_i - \alpha_i u_i$ is nonzero. Here “terminate” means that α_i or β_i equals zero at the current step. Suppose gGKB does not terminate before the k th iteration, i.e., $\alpha_i \beta_i \neq 0$ for $1 \leq i \leq k$. Then the k -step gGKB process generates an M -orthonormal matrix $V_{k+1} = (v_1, \dots, v_{k+1}) \in \mathbb{R}^{n \times (k+1)}$ and a G -orthonormal matrix $U_{k+1} = (u_1, \dots, u_{k+1}) \in \mathbb{R}^{m \times (k+1)}$, satisfying the relations

$$\begin{aligned} (4.8a) \quad & \beta_1 U_{k+1} e_1 = b, \\ (4.8b) \quad & AV_k = U_{k+1} B_k, \\ (4.8c) \quad & M^\dagger A^\top GU_{k+1} = V_k B_k^\top + \alpha_{k+1} v_{k+1} e_{k+1}^\top, \end{aligned}$$

where e_1 and e_{k+1} are the first and $(k+1)$ th columns of I_{k+1} , and

$$(4.9) \quad B_k = \begin{pmatrix} \alpha_1 & & & & \\ \beta_2 & \alpha_2 & & & \\ & \beta_3 & \ddots & & \\ & & \ddots & \alpha_k & \\ & & & \beta_{k+1} & \end{pmatrix} \in \mathbb{R}^{(k+1) \times k}$$

has full column rank. Note that it may occur that $\beta_{k+1} = 0$, which means gGKB terminates just at the k th step, and in this case $v_{k+1} = \mathbf{0}$.

We emphasize that gGKB will eventually terminate in at most $\min\{m, r\}$ steps, since the column rank of U_k or V_k cannot exceed $\min\{m, r\}$. Using the GSVD of $\{A, L\}$, we can give a detailed description of the “terminate step” of gGKB, defined as

$$(4.10) \quad k_t = \min\{k : \alpha_{k+1} \beta_{k+1} = 0\}.$$

For any closed subspace \mathcal{G} of a Hilbert space, denote by $\mathcal{P}_{\mathcal{G}}$ the projection operator onto \mathcal{G} . We have the following result.

THEOREM 4.2. Define the linear operator \mathcal{A} as (3.11) with $M = A^\top A + L^\top L$, where the GSVD of $\{A, L\}$ is as (1.1). Suppose there are l distinct positive c_i in Σ_A with l subspaces $\mathcal{G}_1, \dots, \mathcal{G}_l$ spanned by the corresponding $p_{A,i}$. Then $k_t = s$, the number of nonzero elements in $\{\mathcal{P}_{\mathcal{G}_1} b, \dots, \mathcal{P}_{\mathcal{G}_l} b\}$.

The following lemma is needed to prove this theorem.

LEMMA 4.3. For any square matrix C and a vector v , define the degree of v with respect to C as

$$\deg_C(v) = \min\{k : \exists p \in \mathcal{P}_k \text{ s.t. } p(C)v = \mathbf{0}\},$$

where \mathcal{P}_k is the set of all polynomials with degrees not bigger than k . Then we have

$$(4.11) \quad \deg_{AM^\dagger A^\top}(b) = \deg_{M^\dagger A^\top A}(M^\dagger A^\top b) = s.$$

Proof. First notice that $\deg_C(v)$ is nothing but the maximum rank of $\{C^i v\}_{i=0}^k$ with respect to $k \geq 0$. By Theorem 3.5, we have $AM^\dagger A^\top = A\tilde{X}_1\tilde{X}_1^\top A^\top$. Using the relation $A\tilde{X}_1 = P_A \Sigma_A$, we have $AM^\dagger A^\top = P_A(\Sigma_A \Sigma_A^\top)P_A^\top$, which is the eigenvalue decomposition of $AM^\dagger A^\top$. Thus, the positive eigenvalues of $AM^\dagger A^\top$ are $1, c_{q_1+1}^2, \dots, c_{q_1+q_2}^2$ with the corresponding eigenvectors being the columns of $(P_{A1} \ P_{A2})$, and the corresponding eigenspaces are subspaces $\mathcal{G}_1, \dots, \mathcal{G}_l$. Denote the l distinct positive eigenvalues by $\lambda_1, \dots, \lambda_l$ and let G_i be those matrices with 2-orthonormal columns spanning \mathcal{G}_i for $1 \leq i \leq l$. Then we can write the eigenvalue decomposition of $AM^\dagger A^\top$ as $AM^\dagger A^\top = \sum_{i=1}^l \lambda_i G_i G_i^\top$, and we have $\mathcal{P}_{\mathcal{G}_i} = G_i G_i^\top$. For each $j \geq 0$, it follows that

$$w_j := (AM^\dagger A^\top)^j b = \sum_{i=1}^l (\lambda_i G_i G_i^\top)^j b = \sum_{i=1}^l \lambda_i^j G_i G_i^\top b,$$

since G_i are mutually 2-orthogonal. Without loss of generality, suppose the first s elements in $\{\mathcal{P}_{\mathcal{G}_1} b, \dots, \mathcal{P}_{\mathcal{G}_l} b\}$ are nonzero and $g_i = \mathcal{P}_{\mathcal{G}_i} b / \|\mathcal{P}_{\mathcal{G}_i} b\|_2 \neq \mathbf{0}$ for $1 \leq i \leq s$. Then $\{g_i\}_{i=1}^s$ are mutually 2-orthogonal, and

$$w_j = \sum_{i=1}^s \lambda_i^j g_i \|\mathcal{P}_{\mathcal{G}_i} b\|_2 = \sum_{i=1}^s \lambda_i^j g_i (g_i^\top b),$$

since

$$g_i^\top b = (G_i G_i^\top b)^\top b / \|\mathcal{P}_{\mathcal{G}_i} b\|_2 = \|G_i^\top b\|_2^2 / \|G_i^\top b\|_2 = \|\mathcal{P}_{\mathcal{G}_i} b\|_2.$$

Thus, the rank of $\{w_j\}_{j=0}^k$ is at most s , leading to $\deg_{AM^\dagger A^\top}(b) \leq s$. On the other hand, for $1 \leq k \leq s$, by setting $\bar{w}_j := g_j (g_j^\top b)$, we have $(w_1 \dots, w_k) = (\bar{w}_1, \dots, \bar{w}_s) T_k$, where

$$T_k = \begin{pmatrix} 1 & \lambda_1 & \dots & \lambda_1^{k-1} \\ 1 & \lambda_2 & \dots & \lambda_2^{k-1} \\ \vdots & \vdots & \dots & \vdots \\ 1 & \lambda_s & \dots & \lambda_s^{k-1} \end{pmatrix} =: \begin{pmatrix} T_{k1} \\ T_{k2} \\ \vdots \\ k \end{pmatrix} \begin{matrix} k \\ s-k \end{matrix}.$$

Since $\lambda_i \neq \lambda_j$ for $1 \leq i \neq j \leq k$, the Vandermonde matrix T_{k1} is invertible, and thereby T_k has full column rank. Therefore, the rank of $\{w_i\}_{i=1}^k$ is k for $1 \leq k \leq s$, leading to $\deg_{AM^\dagger A^\top}(b) \geq s$. This proves $\deg_{AM^\dagger A^\top}(b) = s$.

To prove $\text{deg}_{M^\dagger A^\top A}(M^\dagger A^\top b) = s$, it is sufficient to show

$$\text{rank}(\{(M^\dagger A^\top A)^j M^\dagger A^\top b\}_{j=0}^k) = \text{rank}(\{w_j\}_{j=0}^k)$$

for any $k \geq 0$. Notice that

$$(M^\dagger A^\top A)^j M^\dagger A^\top b = M^\dagger A^\top (AM^\dagger A^\top)^j b = M^\dagger A^\top w_j$$

and

$$M^\dagger A^\top = \tilde{X}_1(A\tilde{X}_1)^\top = \tilde{X}_1 \Sigma_A^\top P_A^\top = (X_1 \ X_2 C_{q_2})(P_{A_1} \ P_{A_2})^\top.$$

Let $\tilde{w}_j = (M^\dagger A^\top A)^j M^\dagger A^\top b$. It follows that $\text{rank}(\{\tilde{w}_j\}_{j=0}^k) \leq \text{rank}(\{w_j\}_{j=0}^k)$. To prove the inverse inequality, suppose $\{w_j\}_{j=0}^k$ are linearly independent. We only need to show $\{\tilde{w}_j\}_{j=0}^k$ are linearly independent. If there exist real numbers μ_0, \dots, μ_k such that $\sum_{j=0}^k \mu_j \tilde{w}_j = \mathbf{0}$, then $M^\dagger A^\top Wz = \mathbf{0}$, where $W = (w_0, \dots, w_k)$ has full column rank and $z = (\mu_0, \dots, \mu_k)^\top$. By the expression of $M^\dagger A^\top$, it follows that $Wz \in \mathcal{N}(M^\dagger A^\top) = \mathcal{R}(P_{A_3})$. On the other hand, from $AM^\dagger A^\top = P_A(\Sigma_A \Sigma_A^\top)P_A^\top$ we get $Wz \in \mathcal{R}(W) \subseteq \mathcal{R}(AM^\dagger A^\top) = \mathcal{R}((P_{A_1} \ P_{A_2}))$. Since $\mathcal{R}((P_{A_1} \ P_{A_2})) \cap \mathcal{R}(P_{A_3}) = \{\mathbf{0}\}$, we obtain $Wz = \mathbf{0} \Rightarrow z = \mathbf{0}$, meaning that $\{\tilde{w}_j\}_{j=0}^k$ are independent. This completes the proof. \square

Proof of Theorem 4.2. Suppose gGKB terminates at the k_t th step. By Proposition 4.1, the rank of $\{u_i\}_{i=1}^{k_t}$ is k_t , implying $k_t \leq \text{deg}_{AM^\dagger A^\top}(b) = s$ by Lemma 4.3. Then we show $k_t \geq s$. Notice from the relations (4.1) and (4.2) that

$$\begin{aligned} \alpha_1 \beta_1 v_1 &= M^\dagger A^\top b, \\ \alpha_{i+1} \beta_{i+1} v_{i+1} &= M^\dagger A^\top A v_i - (\alpha_i^2 + \beta_{i+1}^2) v_i - \alpha_i \beta_i v_{i-1} \end{aligned}$$

for $1 \leq i \leq k_t$, where we have used

$$\begin{aligned} M^\dagger A^\top A v_i &= \alpha_i M^\dagger A^\top u_i + \beta_{i+1} M^\dagger A^\top u_{i+1} \\ &= \alpha_i (\alpha_i v_i + \beta_i v_{i+1}) + \beta_{i+1} (\alpha_{i+1} v_{i+1} + \beta_{i+1} v_i). \end{aligned}$$

Therefore, it follows that

$$v_{i+1} = \frac{1}{\alpha_{i+1} \beta_{i+1}} (M^\dagger A^\top A v_i - (\alpha_i^2 + \beta_{i+1}^2) v_i - \alpha_i \beta_i v_{i-1})$$

for $1 \leq i < k_t$. Combining with $v_1 = \frac{1}{\alpha_1 \beta_1} M^\dagger A^\top b$, the above recursion leads to

$$v_{k+1} = \sum_{i=0}^k \xi_i (M^\dagger A^\top A)^i M^\dagger A^\top b, \quad \xi_k = 1 / \prod_{i=1}^{k+1} \alpha_i \beta_i \neq 0$$

for $1 \leq k < k_t$. Since $\mathbf{0} = \alpha_{k_t+1} \beta_{k_t+1} v_{k_t+1}$ is a linear combination of v_{k_t} and v_{k_t-1} with nonzero coefficients, the above identity implies that $\alpha_{k_t+1} \beta_{k_t+1} v_{k_t+1}$ must be a linear combination of $\{(M^\dagger A^\top A)^i M^\dagger A^\top b\}_{i=0}^{k_t}$ with nonzero coefficients, and thereby $\{(M^\dagger A^\top A)^i M^\dagger A^\top b\}_{i=0}^{k_t}$ is linearly dependent. By Lemma 4.3, it follows that $k_t \geq s$. Finally, we obtain $k_t = s$. \square

Just as the standard GKB can be employed to approximate extreme SVD components, we will utilize gGKB to approximate nontrivial extreme GSVD components.

5. GSVD computation by generalized Golub–Kahan bidiagonalization.

We first show that gGKB can be used to approximate the SVE components. Then we use Theorems 3.5 and 3.6 to relate these approximations to the nontrivial GSVD components.

5.1. Computing nontrivial GSVD components by gGKB. Suppose gGKB does not terminate before the k th step. Then the compact-form SVD of B_k can be written as

$$(5.1) \quad B_k = Y_k \Theta_k H_k^\top, \quad \Theta_k = \text{diag}(\theta_1^{(k)}, \dots, \theta_k^{(k)}), \quad \theta_i^{(k)} > \dots > \theta_k^{(k)} > 0,$$

where $Y_k = (y_1^{(k)}, \dots, y_k^{(k)}) \in \mathbb{R}^{(k+1) \times k}$ and $H_k = (h_1^{(k)}, \dots, h_k^{(k)}) \in \mathbb{R}^{k \times k}$ are two 2-orthonormal matrices. The approximation to the SVE triplet $(c_i, p_{A,i}, x_i)$ of \mathcal{A} is defined as $(\bar{c}_i^{(k)}, \bar{p}_{A,i}^{(k)}, \bar{x}_i^{(k)}) := (\theta_i^{(k)}, U_{k+1} y_i^{(k)}, V_k h_i^{(k)})$. To measure the quality of this approximation, we give the following result.

THEOREM 5.1. *The approximate SVE triplet for \mathcal{A} satisfies*

$$(5.2a) \quad \mathcal{A} \bar{x}_i^{(k)} - \bar{c}_i^{(k)} \bar{p}_{A,i}^{(k)} = 0,$$

$$(5.2b) \quad \mathcal{A}^* \bar{p}_{A,i}^{(k)} - \bar{c}_i^{(k)} \bar{x}_i^{(k)} = \alpha_{k+1} v_{k+1} e_{k+1}^\top y_i^{(k)}.$$

Proof. Note that $\mathcal{A}v = Av$. The first relation can be verified using (4.8b):

$$\mathcal{A} \bar{x}_i^{(k)} - \bar{c}_i^{(k)} \bar{p}_{A,i}^{(k)} = AV_k h_i^{(k)} - \theta_i^{(k)} U_{k+1} y_i^{(k)} = U_{k+1} (B_k h_i^{(k)} - \theta_i^{(k)} y_i^{(k)}) = 0.$$

For the second relation, using (3.21), that is, $\mathcal{A}^* u = M^\dagger A^\top u$, we obtain from (4.8c) that

$$\begin{aligned} \mathcal{A}^* \bar{p}_{A,i}^{(k)} - \bar{c}_i^{(k)} \bar{x}_i^{(k)} &= M^\dagger A^\top U_{k+1} y_i^{(k)} - \theta_i^{(k)} V_k h_i^{(k)} \\ &= (V_k B_k^\top + \alpha_{k+1} v_{k+1} e_{k+1}^\top) y_i^{(k)} - \theta_i^{(k)} V_k h_i^{(k)} \\ &= V_k (B_k^\top y_i^{(k)} - \theta_i^{(k)} h_i^{(k)}) + \alpha_{k+1} v_{k+1} e_{k+1}^\top y_i^{(k)} \\ &= \alpha_{k+1} v_{k+1} e_{k+1}^\top y_i^{(k)}. \end{aligned}$$

The proof is completed. \square

Therefore, the triplet $(\bar{c}_i^{(k)}, \bar{p}_{A,i}^{(k)}, \bar{x}_i^{(k)})$ can be accepted as a satisfied SVE triplet at the iteration where $|\alpha_{k+1} v_{k+1} e_{k+1}^\top y_i^{(k)}|$ is sufficiently small. Using the connection between the SVE of \mathcal{A} and the GSVD of $\{A, L\}$ revealed by Theorem 3.5, the tuple $(\bar{c}_i^{(k)}, \bar{s}_i^{(k)}, \bar{p}_{A,i}^{(k)}, \bar{x}_i^{(k)}) := (\theta_i^{(k)}, (1 - (\theta_i^{(k)})^2)^{1/2}, U_{k+1} y_i^{(k)}, V_k h_i^{(k)})$ can be used as a good approximation to a GSVD component. To further measure the quality of this approximation, note from (2.2) that

$$(5.3) \quad s_i^2 A^\top A x_i = c_i^2 L^\top L x_i, \quad 1 \leq i \leq r.$$

This is a well-known basic relation for GSVD, which indicates that the nontrivial generalized eigenvalues of the generalized eigenvalue problem $A^\top A x = \lambda L^\top L x$ are $\{\gamma_i^2\}_{i=1}^r$ and the corresponding generalized eigenvectors are $\{x_i\}_{i=1}^r$ [20, section 8.7]. Now we can give the following result.

THEOREM 5.2. *The above approximate GSVD tuple for $\{A, L\}$ satisfies*

$$(5.4) \quad (\bar{s}_i^{(k)})^2 A^\top A \bar{x}_i^{(k)} - (\bar{c}_i^{(k)})^2 L^\top L \bar{x}_i^{(k)} = \alpha_{k+1} \beta_{k+1} M v_{k+1} e_k^\top h_i^{(k)}.$$

Proof. First notice from (5.3) that

$$\begin{aligned} A^\top Ax_i &= (c_i^2 + s_i^2)A^\top Ax_i = c_i^2(A^\top A + L^\top L)x_i = c_i^2 Mx_i, \\ L^\top Lx_i &= (c_i^2 + s_i^2)L^\top Lx_i = s_i^2(A^\top A + L^\top L)x_i = s_i^2 Mx_i \end{aligned}$$

for $1 \leq i \leq r$. Since $\tilde{X}_1 = (x_1, \dots, x_r)$ is an M -orthonormal basis of $(\mathcal{R}(M), \langle \cdot, \cdot \rangle_M)$, it follows that $A^\top A\mathcal{R}(M) \subseteq \mathcal{R}(M)$ and $L^\top L\mathcal{R}(M) \subseteq \mathcal{R}(M)$. Therefore, we have $A^\top A\bar{x}_i^{(k)}, L^\top L\bar{x}_i^{(k)} \in \mathcal{R}(M)$ due to $\bar{x}_i^{(k)} = V_k h_i^{(k)} \in \mathcal{R}(M)$. By Theorem 5.1, we have

$$\begin{aligned} M^\dagger[(\bar{s}_i^{(k)})^2 A^\top A\bar{x}_i^{(k)} - (\bar{c}_i^{(k)})^2 L^\top L\bar{x}_i^{(k)}] &= M^\dagger[A^\top A\bar{x}_i^{(k)} - (\bar{c}_i^{(k)})^2 M\bar{x}_i^{(k)}] \\ &= \theta_i^{(k)} M^\dagger A^\top U_{k+1} y_i^{(k)} - (\theta_i^{(k)})^2 V_k h_i^{(k)} \\ &= \theta_i^{(k)} (V_k B_k^\top + \alpha_{k+1} v_{k+1} e_{k+1}^\top) y_i^{(k)} - (\theta_i^{(k)})^2 V_k h_i^{(k)} \\ &= \alpha_{k+1} \beta_{k+1} v_{k+1} e_k^\top h_i^{(k)}, \end{aligned}$$

where we have used $B_k^\top y_i^{(k)} = \theta_i^{(k)} h_i^{(k)}$ and $B_k h_i^{(k)} = \theta_i^{(k)} y_i^{(k)}$. Multiplying the above equality by M and using $\mathcal{P}_{\mathcal{R}(M)} = MM^\dagger$, we finally obtain (5.4). \square

Combining Theorems 5.1 and 5.2, it is more proper to use the residual norm

$$(5.5) \quad \|r_i^{(k)}\|_2 := \left(\|A\bar{x}_i^{(k)} - \bar{c}_i^{(k)} \bar{p}_{A,i}^{(k)}\|_2^2 + \|(\bar{s}_i^{(k)})^2 A^\top A\bar{x}_i^{(k)} - (\bar{c}_i^{(k)})^2 L^\top L\bar{x}_i^{(k)}\|_2^2 \right)^{1/2}$$

to measure the quality of the approximate GSVD components of A . Since $\|v_{k+1}\|_M = 1$, it follows from (5.4) that

$$(5.6) \quad \|r_i^{(k)}\|_2 / \|(A^\top, L^\top)^\top\|_2 \leq \alpha_{k+1} \beta_{k+1} |e_k^\top h_i^{(k)}|,$$

because $\|M\|_2^{1/2} = \|(A^\top, L^\top)^\top\|_2$. The easily computed quantity $\alpha_{k+1} \beta_{k+1} |e_k^\top h_i^{(k)}|$ is an upper bound of the scaling-invariant relative residual norm $\|r_i^{(k)}\|_2 / \|(A^\top, L^\top)^\top\|_2$, which can be used in a stopping criterion.

We present the pseudocode of the gGKB-based GSVD computation (computing the GSVD components of A) in Algorithm 5.1. We remark that in order to approximate the GSVD components of L , the gGKB of \mathcal{L} should be used; the spirit is the same as that for \mathcal{A} and we omit it. This process can be computed independently from

Algorithm 5.1. The gGKB-based GSVD computation (gGKB_GSVD).

Input: $A \in \mathbb{R}^{m \times n}$, $L \in \mathbb{R}^{p \times n}$, $\text{tol} > 0$

- 1: Initialize: choose a nonzero $b \in \mathbb{R}^m$; form $M = A^\top A + L^\top L$
- 2: Compute $\beta_1, \alpha_1, u_1, v_1$ by gGKB
- 3: **for** $i = 1, 2, \dots, k$, **do**
- 4: Compute $\beta_{k+1}, \alpha_{k+1}, u_{k+1}, v_{k+1}$ by gGKB; form B_k, U_{k+1} and V_k
- 5: Compute the SVD of B_k as (5.1)
- 6: **if** $\alpha_{k+1} \beta_{k+1} |e_k^\top h_i^{(k)}| < \text{tol}$ **then**
- 7: Compute $(\bar{c}_i^{(k)}, \bar{s}_i^{(k)}, \bar{p}_{A,i}^{(k)}, \bar{x}_i^{(k)}) := (\theta_i^{(k)}, (1 - (\theta_i^{(k)})^2)^{\frac{1}{2}}, U_{k+1} y_i^{(k)}, V_k h_i^{(k)})$
- 8: **end if**
- 9: **end for**

Output: Approximate GSVD components $(\bar{c}_i^{(k)}, \bar{s}_i^{(k)}, \bar{p}_{A,i}^{(k)}, \bar{x}_i^{(k)})$

that of \mathcal{A} , allowing the GSVD components of A and L to be computed in parallel. Furthermore, it is possible to join the two gGKB processes together to save some repeated computations. A systematic comparison of the parallel and joint gGKB processes for large-scale GSVD computations will be explored in future research.

5.2. Convergence and accuracy. We provide preliminary results about the convergence and accuracy of gGKB_GSVD for GSVD computation. The following result demonstrates the good property of gGKB_GSVD at the terminate step.

THEOREM 5.3. *Following the notation and assumptions of Theorem 4.2, then at the k_t th step, the gGKB_GSVD algorithm computes exactly k_t GSVD components corresponding to the nonzero elements in $\{\mathcal{P}_{\mathcal{G}_1}b, \dots, \mathcal{P}_{\mathcal{G}_s}b\}$.*

Proof. By Theorems 5.1 and 5.2, at the terminate step of gGKB, it computes the exact SVE components of \mathcal{A} , which are also the exact GSVD components of $\{A, L\}$ by Theorem 3.5. Following the notation in the proof of Lemma 4.3, we need to prove that the s vectors $\bar{p}_{A,i}^{(s)}$ belong separately to the invariant subspaces $\mathcal{G}_1, \dots, \mathcal{G}_s$. Since $\theta_i^{(s)} > 0$ have different values and \mathcal{G}_i are mutually 2-orthogonal, these $\bar{p}_{A,i}^{(s)}$ must belong to different invariant subspaces. Therefore, we only need to prove $\mathcal{P}_{\mathcal{G}}\bar{p}_{A,i}^{(s)} = \bar{p}_{A,i}^{(s)}$ for each $1 \leq i \leq s$, where $\mathcal{G} = \mathcal{G}_1 \oplus \dots \oplus \mathcal{G}_s$. From the proof of Lemma 4.3, we have $\bar{p}_{A,i}^{(s)} \in \mathcal{K}_s(AM^\dagger A^\top, b) = \text{span}\{w_i\}_{i=0}^{s-1}$, and

$$\widetilde{W}_s := (w_0, \dots, w_{s-1}) = \widetilde{G}_s \begin{pmatrix} g_1^\top b & & & \\ & \ddots & & \\ & & g_1^\top b & \\ & & & \ddots \\ & & & & g_1^\top b \end{pmatrix} \begin{pmatrix} 1 & \lambda_1 & \dots & \lambda_1^{s-1} \\ 1 & \lambda_2 & \dots & \lambda_2^{s-1} \\ \vdots & \vdots & \dots & \vdots \\ 1 & \lambda_s & \dots & \lambda_s^{s-1} \end{pmatrix} =: \widetilde{G}_s \Lambda_s T_s,$$

where $\widetilde{G}_s = (g_1, \dots, g_s)$. Since $g_i^\top b \neq 0$ and T_s is nonsingular, it follows that $\mathcal{R}(\widetilde{W}_s) = \mathcal{R}(\widetilde{G}_s)$. Thus, we can write $\bar{p}_{A,i}^{(s)}$ as $\bar{p}_{A,i}^{(s)} = \widetilde{G}_s z$ with a nonzero $z \in \mathbb{R}^s$. Now we immediately obtain

$$\mathcal{P}_{\mathcal{G}}\bar{p}_{A,i}^{(s)} = \sum_{i=1}^s \mathcal{P}_{\mathcal{G}_i}\bar{p}_{A,i}^{(s)} = \sum_{i=1}^s G_i G_i^\top (\widetilde{G}_s z) = (\widetilde{G}_s \widetilde{G}_s^\top) \widetilde{G}_s z = \bar{p}_{A,i}^{(s)},$$

which is the desired result. \square

To investigate the convergence behavior of the approximations, we give the following result to describe the convergence speed of the Ritz values $\theta_i^{(k)}$.

THEOREM 5.4. *For any $1 \leq i \leq q_1 + q_2$, let*

$$(5.7) \quad \phi_i = \arccos \frac{|c_i p_{A,i}^\top b|}{\|\sum_A^\top P_A^\top b\|_2}.$$

Then at the k th iteration of gGKB_GSVD, it holds for $1 \leq i \leq k$ that

$$(5.8) \quad 0 \leq c_i^2 - (\theta_i^{(k)})^2 \leq (c_1^2 - c_r^2) \left(\frac{\kappa_i^{(k)} \tan \phi_i}{C_{k-i}(1 + 2\gamma_i)} \right),$$

where $C_j(t)$ is the j th Chebyshev polynomial

$$C_j(t) = \frac{1}{2} \left[(t + \sqrt{t^2 - 1})^j + (t - \sqrt{t^2 - 1})^{-j} \right], \quad |t| \geq 1,$$

and

$$\gamma_i = \frac{c_i^2 - c_{i+1}^2}{c_{i+1}^2 - c_r^2}, \quad \kappa_i^{(k)} = \prod_{j=1}^{i-1} \frac{(\theta_j^{(k)})^2 - c_r^2}{(\theta_j^{(k)})^2 - c_i^2} \quad (i > 1), \quad \kappa_1^{(k)} = 1 \quad (i = 1).$$

Proof. Using the relations (4.7) with $G = Z = I_m$ and $W_r = \tilde{X}_1$, it follows that the coordinate representation of **gGKB** is the standard GKB of $A\tilde{X}_1$ with starting vector b . This GKB process is equivalent to the symmetric Lanczos process of $(A\tilde{X}_1)^\top A\tilde{X}_1 \in \mathbb{R}^{r \times r}$ with starting vector $(A\tilde{X}_1)^\top b$, which generates 2-orthogonal vectors $\{u_i\}_{i=1}^k$ and the symmetric tridiagonal matrix $B_k^\top B_k$; see, e.g., [32]. Since $A\tilde{X}_1 = P_A \Sigma_A$, it follows that $(A\tilde{X}_1)^\top A\tilde{X}_1 = I_r (\Sigma_A^\top \Sigma_A) I_r^\top$ is the eigenvalue decomposition of $(A\tilde{X}_1)^\top A\tilde{X}_1$, and $(A\tilde{X}_1)^\top b = \Sigma_A^\top P_A^\top b$. Since the i th eigenvector of $(A\tilde{X}_1)^\top A\tilde{X}_1$ is e_i and

$$(A\tilde{X}_1)^\top b = \Sigma_A^\top P_A^\top b = ((P_{A1}^\top b)^\top \quad (P_{A2}^\top b)^\top \quad \mathbf{0})^\top,$$

the angle between $(A\tilde{X}_1)^\top b$ and e_i for $q_1 + q_2 + 1 \leq i \leq r$ is $\pi/2$, and for $1 \leq i \leq q_1 + q_2$ the angle is expressed as (5.7). Notice that the eigenvalues of $B_k^\top B_k$ are $(\theta_i^{(k)})^2$. Using the convergence theory of the symmetric Lanczos process (see, e.g., [48, Theorem 6.4]), we immediately obtain (5.8). \square

Theorem 5.4 indicates that the convergence rate of $\theta_i^{(k)}$ primarily depends on two factors: the closeness between b and the corresponding vector $p_{A,i}$ and the degree of separation of c_i from others. Therefore, usually we can expect rapid convergence to the extreme and well-separated positive c_i . Note again that the approximations will not converge to the GSVD components corresponding to those zero c_i , since the angle between $(A\tilde{X}_1)^\top b$ and e_i is $\pi/2$ for $q_1 + q_2 + 1 \leq i \leq r$. The convergence behavior of $\tilde{p}_{A,i}^{(k)}$ and $\tilde{x}_i^{(k)}$ can also be described similarly based on the convergence theory of the symmetric Lanczos process, but the mathematical expressions are more complex. Interested readers can refer to [48, section 6.6]

We remark that all the aforementioned results are derived for the **gGKB** with exact computations, i.e., we do not take into account rounding errors and computational errors arising from iteratively solving (4.3). In the presence of rounding errors, the Lanczos-type iterative process behaves very differently from that in exact arithmetic. One well-known result is that the orthogonality of u_i and v_i will be gradually lost, which leads to a delay of convergence of approximations and the appearance of spurious convergent quantities [32]. Also, the inaccurate computation of $M^\dagger \bar{s}$ may affect the final accuracy of the approximations. These issues for **gGKB_GSVD** will be addressed in future work. We will demonstrate several of them in the subsequent numerical experiments.

6. Experimental results. We report some experimental results to demonstrate the performance of **gGKB_GSVD** for computing nontrivial extreme GSVD components. All the experiments are performed in MATLAB R2023b using double precision. The codes are available at https://github.com/Machealb/gsvd_iter. For the starting vector of **gGKB** for A and L , we use the random vector $b = \text{randn}(m, 1)$ and $b = \text{randn}(p, 1)$ with random seed `rng(2024)`, respectively.

Example 1. The matrix pair $\{A, L\}$ is constructed as follows. Set $m = n = p = 1000$. Let $C_A = \text{diag}(\{c_i\}_{i=1}^n)$ with $c(1) = 1, c(2) = 0.95, c(3) = 0.90, c(4 : n - 3) = \text{linspace}(0.88, 0.12, n - 6)$, and $c(n - 2) = 0.1, c(n - 1) = 0.05, c(n) = 0.01$, and let $S_L = \text{diag}(\{s_i\}_{i=1}^n)$ with $s_i = (1 - c_i^2)^{1/2}$. Then let W be an orthogonal matrix by

letting $W = \text{gallery}(\text{'orthog'}, n, 2)$ and $D = \text{diag}(\text{linspace}(1, 100, n))$. Finally, let $A = C_A W^\top D$ and $L = S_L W^\top D$. By construction, $\{A, L\}$ is a regular matrix pair, and the i th generalized singular value of $\{A, L\}$ is c_i/s_i , where the corresponding generalized singular vectors are the i th columns of I_n, I_n , and $D^{-1}W$.

In this experiment, we use `gGKB_GSVD` to compute several largest and smallest generalized singular values and show the convergence behavior of Ritz values $\theta_i^{(k)}$, where `gGKB` is performed with and without reorthogonalization. This is a small-scale problem, and therefore we directly compute M^{-1} for computing $M^{-1}\bar{s}$ at each iteration of `gGKB`. Figure 6.1 shows the convergence of the first three largest and smallest Ritz values, where in the top two subfigures the right vertical lines indicate the values of c_i . There are four findings. (1) If no reorthogonalization is used for `gGKB`, then as the iteration proceeds, the converged Ritz values may suddenly jump up (they also may jump down for converging to those smallest c_i) to become a ghost and then converge to the next larger c_i ; this phenomenon leads to the appearance of spurious copies of computed c_i . (2) If `gGKB` is performed with full reorthogonalization, the convergence of the Ritz values remains regular, and the first three largest and smallest Ritz values converge to the first three largest and smallest c_i , respectively. (3) The final accuracy of the approximated c_i is around $\mathcal{O}(\mathbf{u})$, where $\mathbf{u} = 2^{-53} \approx 10^{-16}$ is the roundoff unit of double precision. (4) The convergence to those largest c_i is faster than the convergence to those smallest c_i ; also, the convergence to c_1, c_2 and c_n, c_{n-1} is faster than that to c_3 and c_{n-2} . This is because c_1, c_2 are more well-separated from others than c_3 ; the same reason applies to c_n, c_{n-1} , and c_{n-2} .




FIG. 6.1. Convergence and accuracy of approximations to c_i , Example 1. Top: convergence of Ritz values $\theta_i^{(k)}$ to largest/smallest c_i by `gGKB_GSVD` without reorthogonalization (left) and with full reorthogonalization (right). Bottom: error curves (full reorthogonalization).




FIG. 6.2. Convergence and accuracy of approximations to s_i , Example 1. Top: convergence of Ritz values $\theta_i^{(k)}$ to largest/smallest s_i by `gGKB_GSVD` without reorthogonalization (left) and with full reorthogonalization (right). Bottom: error curves (full reorthogonalization).

We also test using the `gGKB` of \mathcal{L} to approximate several largest and smallest s_i . The convergence behavior of the Ritz values and error curves are plotted in Figure 6.2. In addition to the findings closely resembling those depicted in Figure 6.1, there are two additional insights. First, we find that the first three smallest Ritz values converge to s_2, s_3, s_4 instead of s_1, s_2, s_3 . This is because $s_1 = 0$, which cannot be converged upon by Ritz values, as revealed by Theorems 5.3 and 5.4. Therefore, we should use `gGKB` of \mathcal{A} to compute the generalized singular values with value ∞ . Second, we find from the bottom two subfigures that those smallest s_i can be approximated more quickly than the largest ones, due to their well-separated locations. Given that these smallest s_i correspond to those largest c_i , it is unsurprising that they exhibit similar convergence behaviors.

Example 2. The matrix A named `well1850` is taken from the SuiteSparse matrix collection [13], and the matrix L is set as

$$L = \begin{pmatrix} 1.1 & -1 & & & \\ & 1.1 & -1 & & \\ & & \ddots & \ddots & \\ & & & 1.1 & -1 \end{pmatrix} \in \mathbb{R}^{(n-1) \times n}.$$

This is a regular matrix pair. We use the MATLAB built-in function `gsvd.m` to compute the full GSVD of $\{A, L\}$ as the baseline of comparison.




FIG. 6.3. Error curves of the approximate GSVD components by `gGKB_GSVD` and relative residual norm with its upper bound, Example 2. Left: approximations to the 1st GSVD components. Right: approximations to the n th GSVD components.

In this experiment, we test the performance of `gGKB_GSVD` for computing the first and n th GSVD components of A for a matrix pair with nonsquare matrices. We only show the results for the `gGKB` of \mathcal{A} and omit the similar results for the `gGKB` of \mathcal{L} . Full reorthogonalization is used and M^{-1} is computed directly. The errors for the approximated generalized singular vectors are measured by $\sin \angle(x, y)$ between two vectors. We also plot the variation of the relative residual norm and its upper bound $\alpha_{k+1}\beta_{k+1}|e_k^\top h_i^{(k)}|$. Figure 6.3 shows that `gGKB_GSVD` can approximate very well the two group extreme GSVD components, with final accuracy around $\mathcal{O}(\mathbf{u})$. The upper bound $\alpha_{k+1}\beta_{k+1}|e_k^\top h_i^{(k)}|$ exhibits a nearly identical decreasing trend as the relative residual norm. Therefore, it is a highly suitable quantity to be employed in the stopping criterion. We also observe that the convergence to the first GSVD components is faster than the convergence to the n th.

Example 3. The matrix pair $\{A, L\}$ is constructed as follows. Set $m = n = p = 1000$ and set $r = 900$. Let $C_A = (\Sigma_A, \mathbf{0})$ with $C_A = \text{diag}(\{c_i\}_{i=1}^r)$, where $c(1) = 0.99, c(2) = 0.98, c(3:r-2) = \text{linspace}(0.96, 0.06, r-4)$, and $c(r-1) = 0.04, c(r) = 0.02$, and let $S_L = (\Sigma_L, \mathbf{0})$ with $S_L = \text{diag}(\{s_i\}_{i=1}^r)$ and $s_i = (1 - c_i^2)^{1/2}$. Then let $W = \text{gallery}(\text{'orthog'}, n, 2)$ be an orthogonal matrix and $D = \text{diag}(\text{linspace}(1, 10, n))$. Finally, let $A = C_A W^\top D$ and $L = S_L W^\top D$. By construction, we have $\text{rank}((A^\top, L^\top)^\top) = r < n$, and the nontrivial GSVD components are c_i, s_i , and the i th columns of I_n, I_n , and $D^{-1}W$ for $1 \leq i \leq r$. For each nontrivial x_i , we compute $\mathcal{P}_{\mathcal{R}(M)} x_i = MM^\dagger x_i$ to get the corresponding right generalized singular vector belonging to $\mathcal{R}(M)$. We use `gGKB_GSVD` to compute x_i that belongs to $\mathcal{R}(M)$.

In this experiment, we test the performance of `gGKB_GSVD` for computing the first and r th GSVD components for a nonregular matrix pair. We directly compute M^\dagger and use full reorthogonalization for `gGKB`. Figure 6.4 shows very good performance of the algorithm: (1) the two extreme GSVD components $(c_i, p_{A,i}, x_i)$ for $i = 1, r$ can be approximated with final accuracy around $\mathcal{O}(\mathbf{u})$; (2) the relative residual norm and its upper bound $\alpha_{k+1}\beta_{k+1}|e_k^\top h_i^{(k)}|$ follow nearly identical decreasing curves, with both eventually stabilizing at a level around $\mathcal{O}(\mathbf{u})$. Again, we observe that the convergence to the first GSVD components is faster than the convergence to the r th.

Example 4. The matrix pair $\{A, L\}$ is constructed as follows. Set $m=n=p=10000$. Let $C_A = \text{diag}(\{c_i\}_{i=1}^n)$ with $c(1) = 0.99, c(2) = 0.97, c(3:n-2) = \text{linspace}(0.95, 0.15, n-4)$, and $c(n-1) = 0.1, c(n) = 0.05$. Let $S_L = \text{diag}(\{s_i\}_{i=1}^n)$ with




FIG. 6.4. Error curves of the approximate GSVD components by `gGKB_GSVD` and relative residual norm with its upper bound, where $\text{rank}((A^\top, L^\top)^\top) = r < n$, Example 3. Left: approximations to the 1st GSVD components. Right: approximations to the r th GSVD components.

$s_i = (1 - c_i^2)^{1/2}$. Then let $W = \text{gallery}(\text{'orthog'}, n, 2)$ be an orthogonal matrix and $D = \text{diag}(\text{linspace}(1, 10, n))$. Finally, let $A = C_A W^\top D$ and $L = S_L W^\top D$. By construction, $\{A, L\}$ is a regular matrix pair, and the i th GSVD components are c_i , s_i , and i th columns of I_n , I_n , and $D^{-1}W$.

We use this experiment to demonstrate the impact of inaccuracy in the computation of $M^\dagger \bar{s}$ on the final accuracy of the approximate GSVD components. We use the MATLAB built-in function `lsqr.m` to solve (4.4) iteratively with stopping tolerance $\text{tol} = 10^{-10}, 10^{-8}$ at each iteration of `gGKB`, respectively. Figure 6.5 shows the decrease of relative errors of the first and n th approximate GSVD components with the two stopping tolerances. We observe that the computational accuracy of $M^\dagger \bar{s}$ significantly affects the final accuracy of both the generalized singular values and vectors. As the computational accuracy deteriorates, so does the final accuracy of the computed GSVD components. Further theoretical investigation into this issue should be conducted in future research.

Example 5. The matrix pair $\{A, L\}$ is constructed as follows. Set $m = n = p = 100000$. Let $C_A = \text{diag}(\{c_i\}_{i=1}^n)$ with $c(1) = 1.0, c(2) = 0.99, c(3 : n - 2) = \text{linspace}(0.98, 0.03, n - 4)$, and $c(n - 1) = 0.02, c(n) = 0.01$. Let $S_L = \text{diag}(\{s_i\}_{i=1}^n)$ with $s_i = (1 - c_i^2)^{1/2}$. Then let $D = \text{diag}(\text{linspace}(1, 50, n))$. Finally, let $A = C_A D$ and $L = S_L D$. By construction, $\{A, L\}$ is a regular matrix pair, and the i th GSVD components are c_i , s_i , and i th columns of I_n , I_n , and D^{-1} .

The aim of this experiment is to test the performance of `gGKB_GSVD` for very large matrix pairs and compare it with `JBD_GSVD`, which is an efficient algorithm for computing extreme GSVD components based on the JBD process [28, 56]. Note that both algorithms are Krylov subspace methods and have nested inner-outer structures. Thus, we use the MATLAB built-in function `lsqr.m` to compute the inner iterations for the two algorithms with stopping tolerance $\text{tol} = 10^{-10}$. The convergence behavior for approximating the largest GSVD components $(c_1, p_{A,1}, x_1)$ and the smallest GSVD components $(c_n, p_{A,n}, x_n)$ is shown in Figure 6.6. For the largest components $(c_1, p_{A,1}, x_1)$, both algorithms can approximate them very quickly, even for this very large matrix pair. For the smallest components $(c_n, p_{A,n}, x_n)$, the two algorithms take many more iterations to converge. Although the specific convergence histories differ, the convergence rates of the algorithms are very similar, likely due to the fact that both algorithms are Krylov subspace methods. We also find that the final accuracy




FIG. 6.5. Accuracy of computed GSVD components by $gGKB_GSVD$, where $s = M^\dagger \bar{s}$ at each $gGKB$ iteration is computed by solving (4.4) using $lsqr.m$ with different stopping tolerance tol , Example 4. Top: $tol=10^{-10}$. Bottom: $tol=10^{-8}$.




FIG. 6.6. Comparison of two algorithms for large-scale GSVD computations, which are based on $gGKB$ and JBD processes, respectively. Top: convergence history for approximating the largest GSVD components. Bottom: convergence history for approximating the smallest GSVD components.

of the two algorithms is similar, with both being influenced by the accuracy of the inner iterations. For large-scale matrices, the primary computational bottleneck of `gGKB_GSVD` lies in the computation of the inner iterations. Future work will focus on theoretically analyzing the required accuracy for these inner iterations and exploring strategies to enhance their computational efficiency.

7. Conclusion and outlook. Based on the theory of SVE of linear compact operators, we have provided a new understanding of the GSVD of $\{A, L\}$ with $A \in \mathbb{R}^{m \times n}$ and $L \in \mathbb{R}^{p \times n}$. By defining the positive semidefinite matrix $M = A^\top A + L^\top L$, we have shown that (1) the trivial GSVD components $\{x_i\}$ form a basis for $\mathcal{N}(M)$ and any nontrivial x_i belongs to the coset $\bar{x}_i + \mathcal{N}(M)$, where $\bar{x}_i \in \mathcal{R}(M)$ is a nontrivial GSVD component; (2) the nontrivial GSVD components of A and L are just the SVEs of the linear operators $\mathcal{A} : (\mathcal{R}(M), \langle \cdot, \cdot \rangle_M) \rightarrow (\mathbb{R}^m, \langle \cdot, \cdot \rangle_2)$, $v \mapsto Av$ and $\mathcal{L} : (\mathcal{R}(M), \langle \cdot, \cdot \rangle_M) \rightarrow (\mathbb{R}^p, \langle \cdot, \cdot \rangle_2)$, $v \mapsto Lv$, respectively. As a direct application of this result, we have developed an operator-type GKB for \mathcal{A} and \mathcal{L} , leading to a novel `gGKB` process. We have used the GSVD of $\{A, L\}$ to study basic properties of `gGKB` and proposed the `gGKB_GSVD` algorithm to compute several nontrivial extreme GSVD components of large-scale matrix pairs. Preliminary results about convergence and accuracy of `gGKB_GSVD` for GSVD computation have been provided, and numerical experiments are presented to demonstrate the effectiveness of this method.

The idea of this paper offers potential directions for developing new algorithms for large-scale GSVD computation. Note that the SVE of \mathcal{A} or \mathcal{L} can be treated as a “weighted” SVD, where the weight matrix M induces a non-Euclidean inner product. Therefore, existing SVD algorithms based on Krylov subspace projection may be modified to approximate the SVE and consequently, the nontrivial GSVD components.

Acknowledgments. The author thanks the two anonymous referees and Editor, Professor Michael Berry, for their detailed reading of the manuscript and many invaluable comments.

REFERENCES

- [1] O. ALTER, P. O. BROWN, AND D. BOTSTEIN, *Generalized singular value decomposition for comparative analysis of genome-scale expression data sets of two different organisms*, Proc. Natl. Acad. Sci. USA, 100 (2003), pp. 3351–3356, <https://doi.org/10.1073/pnas.0530258100>.
- [2] F. ALVARRUIZ, C. CAMPOS, AND J. E. ROMAN, *Thick-restarted joint Lanczos bidiagonalization for the GSVD*, J. Comput. Appl. Math., 440 (2024), 115506, <https://doi.org/10.1016/j.cam.2023.115506>.
- [3] M. ARIOLI, *Generalized Golub–Kahan bidiagonalization and stopping criteria*, SIAM J. Matrix Anal. Appl., 34 (2013), pp. 571–592, <https://doi.org/10.1137/120866543>.
- [4] Z. BAI, J. DEMMEL, J. DONGARRA, A. RUHE, AND H. VAN DER VORST, *Templates for the Solution of Algebraic Eigenvalue Problems: A Practical Guide*, SIAM, Philadelphia, 2000.
- [5] Z. BAI AND J. W. DEMMEL, *Computing the generalized singular value decomposition*, SIAM J. Sci. Comput., 14 (1993), pp. 1464–1486, <https://doi.org/10.1137/0914085>.
- [6] S. J. BENBOW, *Solving generalized least-squares problems with LSQR*, SIAM J. Matrix Anal. Appl., 21 (1999), pp. 166–177, <https://doi.org/10.1137/S0895479897321830>.
- [7] K. BHUYAN, S. SINGH, AND P. BHUYAN, *Application of generalized singular value decomposition to ionospheric tomography*, Ann. Geophys., 22 (2004), pp. 3437–3444.
- [8] Å. BJÖRCK, *Numerical Methods for Least Squares Problems*, SIAM, Philadelphia, 1996.
- [9] N. A. CARUSO AND P. NOVATI, *Convergence analysis of LSQR for compact operator equations*, Linear Algebra Appl., 583 (2019), pp. 146–164, <https://doi.org/10.1016/j.laa.2019.08.024>.
- [10] M. T. CHU, R. E. FUNDERLIC, AND G. H. GOLUB, *On a variational formulation of the generalized singular value decomposition*, SIAM J. Matrix Anal. Appl., 18 (1997), pp. 1082–1092, <https://doi.org/10.1137/S0895479895287079>.

- [11] J. CHUNG AND A. K. SAIBABA, *Generalized hybrid iterative methods for large-scale Bayesian inverse problems*, SIAM J. Sci. Comput., 39 (2017), pp. S24–S46, <https://doi.org/10.1137/16M1081968>.
- [12] J. CHUNG, A. K. SAIBABA, M. BROWN, AND E. WESTMAN, *Efficient generalized Golub–Kahan based methods for dynamic inverse problems*, Inverse Problems, 34 (2018), 024005, <https://doi.org/10.1088/1361-6420/aaa0e1>.
- [13] T. A. DAVIS AND Y. HU, *The university of Florida sparse matrix collection*, ACM Trans. Math. Software, 38 (2011), pp. 1–25, <https://doi.org/10.1145/2049662.2049663>.
- [14] A. DUMITRASC, C. KRUSE, AND U. RUEDE, *Generalized Golub-Kahan Bidiagonalization for Nonsymmetric Saddle Point Systems*, preprint, arXiv:2310.06952, 2023.
- [15] A. EDELMAN AND Y. WANG, *The GSVD: Where are the ellipses?, Matrix trigonometry, and more*, SIAM J. Matrix Anal. Appl., 41 (2020), pp. 1826–1856, <https://doi.org/10.1137/18M1234412>.
- [16] L. ELDÉN, *A weighted pseudoinverse, generalized singular values, and constrained least squares problems*, BIT, 22 (1982), pp. 487–502, <https://doi.org/10.1007/BF01934412>.
- [17] H. W. ENGL, M. HANKE, AND A. NEUBAUER, *Regularization of Inverse Problems*, Kluwer Academic Publishers, Norwell, MA, 2000.
- [18] L. M. EWERBRING AND F. T. LUK, *Canonical correlations and generalized SVD: Applications and new algorithms*, J. Comput. Appl. Math., 27 (1989), pp. 37–52, [https://doi.org/10.1016/0377-0427\(89\)90360-9](https://doi.org/10.1016/0377-0427(89)90360-9).
- [19] G. GOLUB AND W. KAHAN, *Calculating the singular values and pseudo-inverse of a matrix*, J. Soc. Ind. Appl. Math. Ser. B Numer. Anal., 2 (1965), pp. 205–224, <https://doi.org/10.1137/0702016>.
- [20] G. H. GOLUB AND C. F. VAN LOAN, *Matrix Computations*, 4th ed., The Johns Hopkins University Press, MD, Baltimore, 2013.
- [21] P. HANSEN AND M. HANKE, *A Lanczos algorithm for computing the largest quotient singular values in regularization problems*, in SVD and Signal Processing III, M. Moonen and B. D. Moor, eds., Elsevier Science B.V., Amsterdam, 1995, pp. 131–138.
- [22] M. HOCHSTENBACH, *A Jacobi–Davidson type method for the generalized singular value problem*, Linear Algebra Appl., 431 (2009), pp. 471–487, <https://doi.org/10.1016/j.laa.2009.03.003>.
- [23] M. E. HOCHSTENBACH AND Y. NOTAY, *The Jacobi–Davidson method*, GAMM-Mitt., 29 (2006), pp. 368–382, <https://doi.org/10.1002/gamm.201490038>.
- [24] P. HOWLAND, M. JEON, AND H. PARK, *Structure preserving dimension reduction for clustered text data based on the generalized singular value decomposition*, SIAM J. Matrix Anal. Appl., 25 (2003), pp. 165–179, <https://doi.org/10.1137/S0895479801393666>.
- [25] J. HUANG AND Z. JIA, *On choices of formulations of computing the generalized singular value decomposition of a large matrix pair*, Numer. Algorithms, 87 (2021), pp. 689–718, <https://doi.org/10.1007/s11075-020-00984-9>.
- [26] J. HUANG AND Z. JIA, *Two harmonic Jacobi–Davidson methods for computing a partial generalized singular value decomposition of a large matrix pair*, J. Sci. Comput., 93 (2022), 41, <https://doi.org/10.1007/s10915-022-01993-7>.
- [27] Z. JIA AND H. LI, *The joint bidiagonalization process with partial reorthogonalization*, Numer. Algorithms, 88 (2021), pp. 965–992, <https://doi.org/10.1007/s11075-020-01064-8>.
- [28] Z. JIA AND H. LI, *The joint bidiagonalization method for large GSVD computations in finite precision*, SIAM J. Matrix Anal. Appl., 44 (2023), pp. 382–407, <https://doi.org/10.1137/22M1483608>.
- [29] B. KÅGSTRÖM, *The generalized singular value decomposition and the general $(A-\lambda B)$ -problem*, BIT, 24 (1984), pp. 568–583, <https://doi.org/10.1007/BF01934915>.
- [30] R. KRESS, *Linear Integral Equations*, Springer, New York, 2013, <https://doi.org/10.1007/978-1-4614-9593-2>.
- [31] S. KUO, W. YEIH, AND Y. WU, *Applications of the generalized singular-value decomposition method on the eigenproblem using the incomplete boundary element formulation*, J. Sound Vib., 235 (2000), pp. 813–845, <https://doi.org/10.1006/jsvi.2000.2946>.
- [32] R. M. LARSEN, *Lanczos bidiagonalization with partial reorthogonalization*, DAIMI Rep. Ser., 27 (1998).
- [33] H. LI, *Subspace Projection Regularization for Large-Scale Bayesian Linear Inverse Problems*, preprint, arXiv:2310.18618, 2023.
- [34] H. LI, *The joint bidiagonalization of a matrix pair with inaccurate inner iterations*, SIAM J. Matrix Anal. Appl., 45 (2024), pp. 232–259, <https://doi.org/10.1137/22M1541083>.
- [35] H. LI, *A preconditioned Krylov subspace method for linear inverse problems with general-form Tikhonov regularization*, SIAM J. Sci. Comput., 46 (2024), pp. A2607–A2633, <https://doi.org/10.1137/23M1593802>.

- [36] H. LI, J. FENG, AND F. LU, *Scalable Iterative Data-Adaptive RKHS Regularization*, preprint, arXiv:2401.00656, 2024.
- [37] R.-C. LI, *Bounds on perturbations of generalized singular values and of associated subspaces*, SIAM J. Matrix Anal. Appl., 14 (1993), pp. 195–234, <https://doi.org/10.1137/0614017>.
- [38] Y. LIU, X. SHAN, AND M. SHAO, *A Contour Integral-Based Algorithm for Computing Generalized Singular Values*, preprint, arXiv:2401.00121, 2023.
- [39] K. NAKAMURA, K. NAKADAI, AND G. INCE, *Real-time super-resolution sound source localization for robots*, in Proceedings of the IEEE/RSJ International Conference on Intelligent Robots and Systems, IEEE, 2012, pp. 694–699.
- [40] C. C. PAIGE, *A note on a result of Sun Ji-Guang: Sensitivity of the CS and GSV decompositions*, SIAM J. Numer. Anal., 21 (1984), pp. 186–191, <https://doi.org/10.1137/0721013>.
- [41] C. C. PAIGE, *The general linear model and the generalized singular value decomposition*, Linear Algebra Appl., 70 (1985), pp. 269–284, [https://doi.org/10.1016/0024-3795\(85\)90059-X](https://doi.org/10.1016/0024-3795(85)90059-X).
- [42] C. C. PAIGE, *Computing the generalized singular value decomposition*, SIAM J. Sci. Stat. Comput., 7 (1986), pp. 1126–1146, <https://doi.org/10.1137/0907077>.
- [43] C. C. PAIGE AND M. A. SAUNDERS, *Towards a generalized singular value decomposition*, SIAM J. Numer. Anal., 18 (1981), pp. 398–405, <https://doi.org/10.1137/0718026>.
- [44] C. C. PAIGE AND M. A. SAUNDERS, *LSQR: An algorithm for sparse linear equations and sparse least squares*, ACM Trans. Math. Software, 8 (1982), pp. 43–71, <https://doi.org/10.1145/355984.355989>.
- [45] C. H. PARK AND H. PARK, *A relationship between linear discriminant analysis and the generalized minimum squared error solution*, SIAM J. Matrix Anal. Appl., 27 (2005), pp. 474–492, <https://doi.org/10.1137/040607599>.
- [46] P. T. P. TANG AND E. POLIZZI, *FEAST as a subspace iteration eigensolver accelerated by approximate spectral projection*, SIAM J. Matrix Anal. Appl., 35 (2014), pp. 354–390, <https://doi.org/10.1137/13090866X>.
- [47] A. REFAHI SHEIKHANI AND S. KORDROSTAMI, *New iterative methods for generalized singular-value problems*, Math. Sci., 11 (2017), pp. 257–265, <https://doi.org/10.1007/s40096-017-0223-3>.
- [48] Y. SAAD, *Numerical Methods for Large Eigenvalue Problems*, Revised ed., Classics Appl. Math. 66, SIAM, Philadelphia, 2011.
- [49] T. SAKURAI AND H. SUGIURA, *A projection method for generalized eigenvalue problems using numerical integration*, J. Comput. Appl. Math., 159 (2003), pp. 119–128, [https://doi.org/10.1016/S0377-0427\(03\)00565-X](https://doi.org/10.1016/S0377-0427(03)00565-X).
- [50] J. M. SPEISER AND C. VAN LOAN, *Signal processing computations using the generalized singular value decomposition*, in Real-Time Signal Processing VII, SPIE 495, 1984.
- [51] J.-G. SUN, *Perturbation analysis for the generalized singular value problem*, SIAM J. Numer. Anal., 20 (1983), pp. 611–625, <https://doi.org/10.1137/0720041>.
- [52] J.-G. SUN, *Perturbation theorems for generalized singular values*, J. Comput. Math., 1 (1983), pp. 233–242, <https://www.jstor.org/stable/43692323>.
- [53] B. D. SUTTON, *Stable computation of the CS decomposition: Simultaneous bidiagonalization*, SIAM J. Matrix Anal. Appl., 33 (2012), pp. 1–21, <https://doi.org/10.1137/100813002>.
- [54] C. F. VAN LOAN, *Generalizing the singular value decomposition*, SIAM J. Numer. Anal., 13 (1976), pp. 76–83, <https://doi.org/10.1137/0713009>.
- [55] C. F. VAN LOAN, *Computing the CS and the generalized singular value decompositions*, Numer. Math., 46 (1985), pp. 479–491, <https://doi.org/10.1007/BF01389653>.
- [56] H. ZHA, *Computing the generalized singular values/vectors of large sparse or structured matrix pairs*, Numer. Math., 72 (1996), pp. 391–417, <https://doi.org/10.1007/s002110050175>.